

## Technical Factors Affecting to Production of Sweet Potato Wine

Nguyen Phuoc Minh\*

Tra Vinh University, Vietnam

\*Corresponding Author E-mail: [dr.nguyenphuocminh@gmail.com](mailto:dr.nguyenphuocminh@gmail.com)

### ABSTRACT

*Wine is a popular alcoholic drink. It is prepared from various fruit juices by fermentation. The sweet potato (*Ipomoea batatas*) is a root vegetable with high starch content. The nutritional composition of sweet potato which are important in meeting human nutritional needs including carbohydrates, fibres, carotenes, thiamine, riboflavin, niacin, potassium, zinc, calcium, iron, vitamins A and C and high quality protein. It has a wide range of uses, including food, beverages, medicines, ceremonial and household objects, fishing bait, and animal feed. In this study, we examine the initial moisture content in sweet potato 77.4%, fermentation temperature 28°C in 120 hours. Ratio of yeast supplementation is 20%, gelatin 1% and sweet potato fluid 30% for wine fermentation.*

**Keywords:** *sweet potato, wine, yeast, fermentation*

### INTRODUCTION

Wine is the product of complex interactions between fungi, yeasts and bacteria that commence in the vineyard and continue throughout the fermentation process until packaging. The sweet potato (*Ipomoea batatas* L.) is a dicotyledon plant that belongs to the family Convolvulaceae. Sweet potato has large, starchy and sweet tasting tuberous roots. The roots are rich in starch, sugars, vitamin C, provitamin A, iron and minerals. Some varieties contain colored pigments such as  $\beta$ -carotene and anthocyanin<sup>2,9</sup>. These pigments have antioxidant properties possessing health promoting attributes such as ability to fight cancer, protect against night blindness, delay aging and prevent liver injury. The roots are consumed as fresh vegetable or processed by steaming or boiling, baking, frying, roasting, making chips, converting into flour, etc. Roots of some of the cultivars are used for the preparation of beverage, paste, powder, alcohol drink and natural color<sup>3</sup>. Reports on the production of non-alcoholic beverage from sweet potato are available in the literature<sup>1,8</sup>. An investigation was undertaken to identify the best fermentation process parameters for the production of wine from reconstituted sweet potato (*Ipomoea batatas*) juice and study the effect of storage period and temperature on sensory and optical attributes of wine<sup>7</sup>. Ramesh C. Ray *et al.*<sup>6</sup> examined the proximate composition and sensory evaluation of anthocyanin-rich purple sweet potato (*Ipomoea batatas* L.) wine. Karthik Raja. G *et al.*<sup>4</sup> conducted the production of wine using sweet potato and also to find an effective combinational resource. Initially, white wine was produced using sweet potato. Through a detailed analysis, it was confirmed that the rate of white wine production was higher in sweet potato when compare to grapes and figs. M. O. Oke and T. S. Workneh reviewed sweet potato postharvest processing and preservation technology.

The main purpose of this our research is to investigate the effect of yeast supplementation and fermentation time to the pH change, yeast density and starch residue and ethanol formation.

**MATERIAL AND METHOD**

**Material**

**Fig. 1: Sweet potato**



Sweet potato is supplied from the Mekong River Delta, Vietnam. Yeast *Saccharomyces cerevisiae* is originated from Pasteur Institute, HCMC, Vietnam

**Research method**

**Experiment #1: Effect of the yeast ratio, fermentation time to pH change, yeast density, starch residue, and ethanol formation**

Moisture content in sweet potato before fermentation is 77.4%. We examine different yeast ratios: 17%, 20%, 23%, 26% fermented at the fixed temperature. Then we determine pH, yeast density, starch residue and ethanol formation

**Experiment #2: Effect of sweet potato ratio to wine sensory characteristics**

Gelatin 1%. Weigh 50 gram of sweet potato powder + 180 ml of warm water to dissolve it. The ratio of weet potato is 20%, 30%, 40%, 50%. This milk is mixed with flavour and then homogenized to get the pleasant flavour and aroma.


**Statistical analysis**

All data are processed by Excell.

**RESULT AND DISCUSSION**


**Effect of the yeast ratio, fermentation time to pH change, yeast density, starch residue, and ethanol formation**

**Fig. 2: Change of pH values**




During fermentation, ethanol is insignificantly difference by pH .

Fig. 3: Yeast cell during fermentation




We decide to choose sample with 20% yeast to ferment the sweet potato juice.

Fig. 4: Ethanol formation during fermentation



We decide to choose sample with 20% yeast because it produces more thanol and economic.

Fig. 5: Starch residue by fermentation time



We decide to choose sample with 20% yeast

**Effect of the ratio of the milk and sweet potato wine to product sensory characteristics**

The sensory evaluating board includes 10 members. Each specialist receives 25 ml of the sweet potato wine in glass beaker at the 22°C. We choose sample with 30% milk with 1% gelatin and 69% sweet potato wine to get the best wine quality.

**CONCLUSION**

Sweet potatoes (*Ipomoea batatas* L.) are rich in dietary fibre, minerals, vitamins and antioxidants, such as phenolic acids, anthocyanins, tocopherol and b-carotene. We have successfully investigated the effect of the yeast ratio, fermentation time to pH change, yeast density, starch residue, and ethanol formation during the sweet potato wine fermentation.

**REFERENCES**

1. Coggins, P.C., Kelly, R.A. and Wilbourn, J.A. (2003). Juice yield of sweet potato culls. Session 104C, Fruit and Vegetable Products: Vegetables (Processed), 2003 IFT Annual Meeting. Chicago, USA.
2. Hou, W.C. Chen, Y.C. and Chen, H.J. Antioxidant activities of trypsin inhibitor, a 33 KDa root storage protein of sweet potato (*Ipomoea batatas* (L.) Lam cv. Tainong 57. *Journal of Agricultural and Food Chemistry*, **49(6)**: 2978-2981 (2001)
3. Islam, M.S. and Jalaluddin, M. Sweet potato - A potential nutritionally rich multifunctional food crop for Arkansas. *Journal of Arkansas Agriculture and Rural Development* **4**: 3-7 (2004)
4. Karthik Raja, G. Ezhilarasan, V. Yazhini, K.A. Sridhar, S. And Chinnathambi, V. Comparative evaluation of white wine production from different carbohydrate rich substrates using air-lift bioreactor. *International Journal of Pharma and Bio Sciences*, **3(1)**: 392-404 (2012)
5. M. O. Oke and T. S. Workneh (2013). A review on sweet potato postharvest processing and preservation technology. *African Journal of Agricultural Research*, 8(40): 4990-5003.
6. Ramesh C. Ray, Sandeep K. Panda, Manas R. Swain & P. Sethuraman Sivakumar, Proximate composition and sensory evaluation of anthocyanin-rich purple sweet potato (*Ipomoea batatas* L.) wine. *International Journal of Food Science and Technology*, **1**: 1-7 (2011)
7. Sanjib Kumar Paul, Himjyoti Dutta, Charu Lata Mahanta and Prasanna Kumar, G.V. Process standardization, characterization and storage study of a sweet potato (*Ipomoea batatas* L.) wine. *International Food Research Journal*, **21(3)**: 1149-1156 (2014)
8. Wireko-Manu, F.D. Ellis, W.O. and Oduro, I. Production of a non-alcoholic beverage from sweet potato (*Ipomoea batatas* L.). *African Journal of Food Science*, **4(4)**: 180-183 (2010)
9. Yamakawa, O. (1998). Development of new cultivation and utilization system for sweet potato towards the 21st century. Proceedings of International workshop on sweet potato production system towards the 21st century, pp.1-8.