

## Growth, Yield Attributes and Yield of Ajwain (*Trachyspermum ammi*) Influenced by Integrated Weed Management

N. Nalini, A. Srinivas\*, T. Ramprakash and V. Praveen Rao

College of Agriculture, Professor Jaya Shankar Telangana State Agricultural University  
Hyderabad, Telangana, India

\*Corresponding Author E-mail: [nalinitara@gmail.com](mailto:nalinitara@gmail.com)

Received: 21.07.2017 | Revised: 4.08.2017 | Accepted: 5.08.2017

### ABSTRACT

An investigation on “Studies on Integrated Weed Management in Ajwain (*Trachyspermum ammi*)” was carried out during rabi 2012 at College farm, Rajendranagar, Hyderabad. Hand weeding at 20, 40 and 60 DAS recorded significantly higher plant height and dry matter production at harvest of the crop, yield attributes and yield over the other treatments followed by application of oxyfluorfen @ 0.12 kg a.i ha<sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg a.i ha<sup>-1</sup> as PoEat 20 DAS.

**Key words:** Ajwain, Herbicides, Weed Management, Yield, Economics

### INTRODUCTION

*Trachyspermum ammi* misprauge, commonly known as ajwain or bishop’s weed and vaamu in telugu. *Trachyspermum ammi* is a native of Egypt and is cultivated in Iraq, Iran, Afghanistan, Pakistan and India. In India, it is cultivated in Andhra Pradesh, Madhya Pradesh, Uttar Pradesh, Gujarat, Rajasthan, Maharashtra, Bihar and West Bengal. *Trachyspermum ammi* sprauge belonging to family Apiaceae is a highly valued medicinally important seed spice. Ajwain is a profusely branched annual herb of 60-120 cm tall. Ajwain is traditionally arabi season crop and its productivity is low due to several factors and one of them is uncontrolled weed growth during the critical periods, and also at subsequent stages of the crop growth. Ajwain is generally infested with grasses, sedges and

broad leaved weeds which smother ajwain at early stages of crop growth ultimately cause yield loss. Among the several weed control measures hand weeding is mostly practiced. Of late this practice has become uneconomical due to increased cost of manual labour, besides non availability of labour during peak periods of agricultural operations and time taken for weeding make the practice of hand weeding is not possible always. National Research Centre on Seed Spices (NRCSS)<sup>5</sup> recommends application of pendimethalin @ 1.0 kg a.i ha<sup>-1</sup> as pre-emergence for weed management in ajwain. However, the chemical is costly and not cost effective to farmers. In the current situations, when labour availability is serious problem, weed management in a crop of 150-180 days duration without post-emergence herbicides is difficult.

**Cite this article:** Nalini, N., Srinivas, A., Ramprakash, T. and Rao, V.P., Growth, Yield Attributes and Yield of Ajwain (*Trachyspermum ammi*) Influenced by Integrated Weed Management, *Int. J. Pure App. Biosci.* 5(4): 925-931 (2017). doi: <http://dx.doi.org/10.18782/2320-7051.5639>

Hence there is a clear need to evaluate the efficacy and toxicity of pre-emergence, post-emergence herbicides and their combinations on ajwain and also the effectiveness of integrated weed management approaches involving the mechanical methods conjunction with pre-emergence herbicide usage. This will lead to identification of effective and economical weed control strategy for productivity enhancement in ajwain.

### MATERIALS AND METHODS

The experiment was carried out at College of Agriculture, Rajendranagar, Hyderabad (Telangana) during the *rabi* season of 2012-13. The experiment comprised of 13 treatments namely, T<sub>1</sub> Pendimethalin @ 1.0 kg *a.i ha*<sup>-1</sup> as PE fb hand weeding at 40 DAS, T<sub>2</sub> - Oxyfluorfen @ 0.12 kg *a.i ha*<sup>-1</sup> as PE fb hand weeding at 40 DAS, T<sub>3</sub> - Pretilachlor @ 0.5 kg *a.i ha*<sup>-1</sup> as PE fb hand weeding at 40 DAS, T<sub>4</sub> - Quizalofop -p-ethyl @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>5</sub> Propanil @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>6</sub> - Pendimethalin @ 1.0 kg *a.i ha*<sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>7</sub> Oxyfluorfen @ 0.12 kg *a.i ha*<sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>8</sub> - Pretilachlor @ 0.5 kg *a.i ha*<sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>9</sub> - Pendimethalin @ 1.0 kg *a.i ha*<sup>-1</sup> as PE fb propanil @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>10</sub> - Oxyfluorfen @ 0.12 kg *a.i ha*<sup>-1</sup> as PE fb propanil @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>11</sub> - Pretilachlor @ 0.5 kg *a.i ha*<sup>-1</sup> as PE fb propanil @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS, T<sub>12</sub> - Hand weeding at 20, 40 and 60 DAS and T<sub>13</sub> - Weedy check. The experiment was laid out in a randomized block design with three replications. The soil of the experimental site was sandy loam with a pH of 7.8 and having 0.35% organic carbon and 226, 18 and 236 kg *ha*<sup>-1</sup> low in available nitrogen and available phosphorous and

medium in potassium, respectively. Irrigation and other cultural practices were adopted as per recommendation. Observations on growth, yield attributing characters, yield and economics were taken. Net returns and benefit: cost (B : C) ratio were also worked out. Weed control efficiency (WCE) was calculated as per the formula suggested by Patil & Patil<sup>6</sup>. Observations on weed dry matter were recorded for whole plot of 4.5 m x 4.0 m, which was used to calculate the WCE. WCE (%) = [(DMC-DMT)/DMC] x 100 where, DMC is the dry matter weight of weeds in control plot and DMT is the dry matter weight of weeds in treated plot. Weed index (WI) was calculated as per the formula suggested by Gill & Kumar<sup>2</sup>. WI (%) = [(X-Y)/X] x 100 where, X is the yield from weed-free plot and Y is the yield from treated plot.

### RESULTS AND DISCUSSION

#### Weed parameters

The weed flora associated with experimental crop consisted of grasses viz., *Cynodon dactylon*, *Dactyloctenium aegyptium* and *Celotia argentia*, sedges viz., *Cyperus rotundus* and broadleaved weeds viz., *Digera arvensis*, *Trianthema portulacastrum*, *Commelina benghalensis*, *Parthenium hysterophorus*, *Euphorbia hirta* and *Hemidismus indica*. Among all weed spp, *Cynodon dactylon*, *Cyperus rotundus* and *Parthenium hysterophorus* were the most dominant ones. Population of broad leaf weeds at the harvesting stage was missing on account of shorter life span than that of ajwain crop. At harvest the lowest weed dry matter was recorded in hand weeding at 20, 40 and 60 DAS (5.87 g *m*<sup>-2</sup>) which was significantly superior over all other treatments. The next best treatment was oxyfluorfen @ 0.12 kg *a.i ha*<sup>-1</sup> as PE fb quizalofop-p-ethyl @ 0.05 kg *a.i ha*<sup>-1</sup> as PoE at 20 DAS (7.00 g *m*<sup>-2</sup>) and it was on par with oxyfluorfen @ 0.12 kg *a.i ha*<sup>-1</sup> as

PE fb hand weeding at 40 DAS ( $7.06 \text{ g m}^{-2}$ ). Significantly the highest weed dry matter was recorded with weedy check ( $12.91 \text{ g m}^{-2}$ ) shows in Table 1. The studies signified the importance of hand weeding at 40 DAS or application of post emergence herbicides which could benefit the crops in reducing the weed dry matter ultimately increase the crop yields. Similar results were obtained with hand weeding in cumin by Chaudhary and Gupta<sup>1</sup>. Hand weeding at 20, 40 and 60 DAS recorded the highest weed control efficiency of 79.78% followed by application of oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb quizalofop-p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS (72.68%), oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb hand weeding at 40 DAS (70.51%). Significantly lower weed control efficiency was recorded with propaquizafop @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS (37.31%) and quizalofop-p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS (44.63%) because of reduced action of post emergency application of herbicides alone. Maximum weed control efficiency recorded in hand weeding is due to continuous removal of weeds upto 60 days after sowing followed by application of different herbicides. These results are in accordance with the results indicated by Sharma and Jain<sup>3</sup> and Mehriya<sup>4</sup>.

### Growth

The plant height at harvest was significantly higher with hand weeding at 20, 40 and 60 DAS (123 cm) than all other treatments but it was on par with application of oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb quizalofop-p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS (117 cm) and oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb hand weeding at 40 DAS (116 cm). The next best treatment to recorded plant height was pretilachlor @  $0.5 \text{ kg a.i ha}^{-1}$  as PE fb quizalofop-p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE (113 cm), pendimethalin @  $1.0 \text{ kg a.i ha}^{-1}$  fb hand weeding at 40 DAS (112 cm) and

pendimethalin @  $1.0 \text{ kg a.i ha}^{-1}$  fb quizalofop-p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS (107 cm).

At harvest, significantly higher dry matter ( $7386 \text{ kg ha}^{-1}$ ) was recorded with hand weeding at 20, 40 and 60 DAS over all other treatments and it was on par with application of oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb quizalofop-p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS ( $7025 \text{ kg ha}^{-1}$ ), oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb hand weeding at 40 DAS ( $6856 \text{ kg ha}^{-1}$ ). The dry matter was significantly lower in unweeded control ( $4126 \text{ kg ha}^{-1}$ ) than rest of the treatments. The higher dry matter where in hand weeding and post emergence treatments were imposed might be due to checking of weed growth efficiently besides increased availability of nutrients to crop. Similar results were found by Susila and Rajkumar<sup>8</sup>.

### Yield parameters and Yield

Hand weeding gave significantly number of umbels plant<sup>-1</sup>, seeds umbel<sup>-1</sup> and test weight (Table 2) and higher seed yield ( $1,155 \text{ kg ha}^{-1}$ ) and haulm yield ( $1316 \text{ kg ha}^{-1}$ ) over all other treatments (Table 3). Among integrated weed control treatments, oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb quizalofop -p-ethyl @  $0.05 \text{ kg a.i ha}^{-1}$  as PoE at 20 DAS proved efficient in recording higher weed control efficiency (72.68%), seed yield ( $1,019 \text{ kg ha}^{-1}$ ) and haulm yield ( $1221 \text{ kg ha}^{-1}$ ) with better weed index (11.77). This was closely followed by oxyfluorfen @  $0.12 \text{ kg a.i ha}^{-1}$  as PE fb hand weeding at 40 DAS which recorded a seed yield of  $959 \text{ kg ha}^{-1}$  and haulm yield ( $1212 \text{ kg ha}^{-1}$ ) with a weed index of 16.97. These findings are conformity with Meena and Mehta<sup>3</sup> reported that hand weeding and application of pre emergence reduced the dry matter of weeds and thus increased in yield attributing and seed yield of seed spices.

**Table 1: Effect of different integrated weed control treatments on weed dry matter at harvest, WCE (%) and Weed index in ajwain at harvest during rabi, 2012-13**

S.No	Treatments	Weed dry matter (g m <sup>-2</sup> )	WCE (%)	WI(%)
T <sub>1</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	8.45 (70.53)	57.89	35.93
T <sub>2</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	7.06 (48.92)	70.51	16.97
T <sub>3</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	8.81 (76.66)	53.79	45.63
T <sub>4</sub>	Quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	9.63 (91.86)	44.63	54.98
T <sub>5</sub>	Propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	10.24 (104.00)	37.31	57.14
T <sub>6</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	8.63 (73.73)	55.56	33.16
T <sub>7</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	6.80 (45.13)	72.68	11.77
T <sub>8</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	9.31 (85.80)	48.28	48.23
T <sub>9</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	9.10 (81.86)	50.65	42.94
T <sub>10</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	8.87 (77.73)	53.15	36.28
T <sub>11</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	9.31 (85.73)	48.32	53.59
T <sub>12</sub>	Hand weeding at 20, 40 and 60 DAS	5.87 (33.60)	79.75	-
T <sub>13</sub>	Weedy check	12.91 (165.90)	-	64.33
	SEm±	0.37	-	
	CD(0.05)	1.08	-	

Original values are given in parentheses, which were transformed to  $\sqrt{x+1}$

**Table 2: Effect of different integrated weed control treatments on plant height (cm) and dry matter (kg ha<sup>-1</sup>) of ajwain at harvest during rabi, 2012-13**

S.No	Treatments	Plant height(cm)	Drymatter (kg /ha)
T <sub>1</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	112	6217
T <sub>2</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	116	6856
T <sub>3</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	105	6126
T <sub>4</sub>	Quizalfop -p-ethyl @0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	105	5868
T <sub>5</sub>	Propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	104	5669
T <sub>6</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	107	6312
T <sub>7</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb quizalfop-p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	117	7025
T <sub>8</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	113	5640
T <sub>9</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	108	5463
T <sub>10</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	109	6018
T <sub>11</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	103	5166
T <sub>12</sub>	Hand weeding at 20, 40 and 60 DAS	123	7386
T <sub>13</sub>	Weedy check	99	4126
	SEm±	3.1	208
	CD(P=0.05)	8.6	608

**Table 3: Effect of different integrated weed control treatments on yield parameters in ajwain at harvest during rabi, 2012-13**

S.No	Treatments	Umbel/ plant	Seed/ Umbel	1000 seed weight
T <sub>1</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	150	162	1.16
T <sub>2</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	243	195	1.40
T <sub>3</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb hand weeding at 40 DAS	113	118	1.06
T <sub>4</sub>	Quizalfop -p-ethyl @0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	93	89	1.06
T <sub>5</sub>	Propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	82	82	1.00
T <sub>6</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	180	180	1.30
T <sub>7</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb quizalfop-p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	270	203	1.53
T <sub>8</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb quizalfop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	106	103	1.23
T <sub>9</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	120	127	1.16
T <sub>10</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	140	142	1.16
T <sub>11</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	102	94	1.10
T <sub>12</sub>	Hand weeding at 20, 40 and 60 DAS	343	218	1.60
T <sub>13</sub>	Weedy check	64.3	73.3	1.06
	SEm±	6.44	1.81	-
	CD(P=0.05)	18.9	5.31	-

**Table 4: Seed yield and haulm yield of different integrated weed control treatments in ajwain during rabi, 2012-13**

S.No	Treatments	Seed yield (kg ha <sup>-1</sup> )	Haulm yield (kg ha <sup>-1</sup> )
T <sub>1</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb Hand weeding at 40 DAS	740	886
T <sub>2</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb Hand weeding at 40 DAS	959	1212
T <sub>3</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb Hand weeding at 40 DAS	628	834
T <sub>4</sub>	Quizalofop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	520	641
T <sub>5</sub>	Propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE at 20 DAS	495	626
T <sub>6</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE	772	931
T <sub>7</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE	1065	1222
T <sub>8</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb quizalofop -p-ethyl @ 0.05 kg a.i ha <sup>-1</sup> as PoE	598	665
T <sub>9</sub>	Pendimethalin @ 1.0 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE	659	787
T <sub>10</sub>	Oxyfluorfen @ 0.12 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE	736	813
T <sub>11</sub>	Pretilachlor @ 0.5 kg a.i ha <sup>-1</sup> as PE fb propaquizafop @ 0.05 kg a.i ha <sup>-1</sup> as PoE	536	673
T <sub>12</sub>	Hand weeding at 20, 40 and 60 DAS	1155	1316
T <sub>13</sub>	Weedy check	412	600
	SEm±	29	31.2
	CD(P=0.05)	84	91

### REFERENCES

- Chaudhary, G. R and Gupta, O. P., Response of cumin (*Cuminumcyminum* L.) to nitrogen application, weed control and sowing methods. *I. J. of Agronomy*. **36**: 212-216 (1991).
- Gill, G.S and Vijay Kumar, K. "Weed index" A new method for reporting weed control trails. *Indian Journal of Agronomy*. **14**: 96-98 (1969).
- Meena, S.S and Mehta, R.S. Effect of weed management on weed indices, yield and economics of fennel (*Foeniculum vulgare*). *Indian Journal of Weed Science*. **41(3&4)**: 195-198 (2009).
- Mehriya, M.L., Yadav, R.S., Jangir, R.P and Poonia, B.L., Nutrient utilization by cumin (*Cuminumcyminum* L.) and weeds as influenced by different weed-control methods. *Indian Journal of Agronomy*. **52**: 1-4 (2007).
- National Research Centre on Seed Spices, Ajmer. (Rajasthan).
- Patil, V. C and Patil, S .V. Studies on weed control in Bamboo. *Indian Journal of Weed Science*. **15 (3)**: 83-86(1983).

7. Sharma, O. L. and Jain, N. K., Effect of levels and time of application of herbicides on seed yield of cumin (*Cuminumcyminum* L.).*Indian Journal of Agricultural Science*.**75 (12):** 812-813 (2005).
8. Susila, T and Rajkumar M. Effect of date of sowing of ajwainsprague on seed yield southern telengana, Andhra Pradesh. *Madras Agricultural Journal*. **98 (1-3):** 39-40(2011).