

Factors Affecting Adherence to Healthy Lifestyle

Bhawana Sharma^{1*} and Mukta Agrawal²

¹Research Scholar, ²Associate Professor

Department of Home Science, University of Rajasthan, Jaipur, Rajasthan

*Corresponding Author E-mail: sharma10bhawana@gmail.com

Received: 15.07.2017 | Revised: 26.07.2017 | Accepted: 27.07.2017

ABSTRACT

The aim of study was to determine reasons for poor adherence to lifestyle recommendations. The objective was to determine: reason for poor adherence to dietary regimen, exercise recommendations, medications, and smoking cessation. Factors affecting adherence to lifestyle changes were categorized into four major categories i.e. socio-demographic factors, cognitive factors, interpersonal factors, unintentional/others. It was found that the major factors responsible for non-adherence were busy schedule, low socioeconomic status, low education level, beliefs, health condition, severe weather condition, cost of medications, side effects of medications, poor memory, lack of motivation, lack of social and family support, unwillingness, tendency to eat out, increasing number of fast food outlet, frequent social gatherings, depression, stress, smoking, patient-physician communication, trust in health-care provider, insufficient information on benefits and use of prescribed medicines and proposed lifestyle changes. No particular intervention strategy can improve the adherence of all patients, research studies reach a conclusion that effective attempts to improve patient adherence depend upon a set of significant factors. Mutual cooperation fosters greater patient satisfaction, lessens the risks of non-adherence, and improves patients' health and fitness.

Key words: Poor adherence, Lifestyle changes, Cognitive factors, Cardiovascular risk.

INTRODUCTION

Cardiovascular diseases and its associated risk factors including hypertension, dyslipidemia, type 2 diabetes, and obesity can be prevented and managed through a combination of healthy lifestyle and medications. Numerous studies have shown the benefit of healthy dietary habits and regular exercise in the prevention and management of type 2 diabetes^{1,2,3,4,5} and obesity^{6,7}. Along with identifying the most

appropriate treatment regimen for an individual with or at high risk of cardiovascular disease, the patient's adherence to therapy is vital in realizing optimal cardiovascular risk reduction. Targeting lifestyle modification amongst patients with lifestyle diseases is effective if the health-care practitioner understands patients' reasons for non-adherence to healthy diet and exercise recommendations.

Cite this article: Sharma, B. and Agrawal, M., Factors Affecting Adherence to Healthy Lifestyle, *Int. J. Pure App. Biosci.* 5(4): 105-116 (2017). doi: <http://dx.doi.org/10.18782/2320-7051.5342>

Despite decades of research, non-adherence to doctors' recommendations remains a major health-care issue⁸. The adherence rate ranges between 20% and 80% depending on the population being studied, used adherence measures, nature of the person's health conditions, and characteristics of the treatment procedures^{9,10}. On average, researchers estimate that 50% of patients do not adhere to medication regimens prescribed by their doctors¹¹ and the rate of nonadherence to lifestyle change recommendations is even higher^{11,12}. Regimen adherence problems are common in individuals with diabetes, thus making glycaemic control difficult to attain^{13,14,15}.

Poor adherence to lifestyle recommendations leads to poor control of the conditions, reduces the effectiveness of therapy and increases the risk of cardiovascular events^{16,17}, lead to relapses, hospitalization, complications, or even death^{8,16,18}. It can also reduce patients' quality of life and, at the broader level, lead to poorer population health outcomes and increased health care costs¹⁹. In addition, higher hospital admission rates resulting from non-adherence increase the cost of medical care^{18,20}. Finally, participants failing to adhere to treatment regimens compromise the results of clinical trials²¹. Poor adherence seems to be a significant barrier to attainment of positive clinical outcome among patient of type 2 diabetes²². Overall, non-adherence to recommended lifestyle and medications represents an important and widespread issue. Supporting adherence to long-term medicines and lifestyle modification is therefore an essential component of patient management. In order to reduce the negative impact of non-adherence, we need to understand the reasons why some patients don't follow their health care provider's recommendations²³. Thus the first step in helping patients with or at risk of cardiovascular disease is determining the factors that could influence the patients' adherence.

The reason of non-adherence to one intervention may be different from another.

This review alluded reasons for non-adherence to recommendations on lifestyle relating to diet, exercise, medications and smoking cessation.

Factors affecting adherence to healthy lifestyle:

I. Socio-demographic factors

1. Age: Studies have shown that old age was associated with better adherence to dietary recommendations^{24,25,26,27,28,29} and smoking cessation²⁸ and overall lifestyle changes²⁹. Elderly people may concentrate more on healthy lifestyle as they are free from family responsibilities and work responsibilities. Another reason for adherence may be the consciousness about health, and desire to improve physical fitness.

Few studies have also shown that old age was found to be associated with non-adherence to physical activity³⁰, dietary recommendations^{31,32} and overall lifestyle changes^{33,34}. It may be due to the health problems, fear of injury as well as attitudinal barriers such as misconceptions about exercise and perceived lack of ability, and dependency on other family members.

Researchers also indicated a U-shape relationship between age and overall lifestyle changes in diabetics³⁵. It was also found in a study that there was no significant relationship of age with dietary, physical activity and medical adherence in hypertensive individuals²⁸.

2. Gender: Women are more weight concern and have self-perception of their body. Men's attitude towards food is generally uncomplicated and enjoyable though they are more frequently overweight and have higher risk of cardiovascular diseases. In several studies females have been reported to be more likely than males to mention healthy dietary habits^{36,37,38,39,40}.

3. Socio-economic status: Socio-economic status was found to be positively related with adherence to dietary²⁸, physical activity³¹ and medical recommendations²². Income was also found to be positively related with adherence to dietary, physical activity and medical recommendation³¹. It was also revealed in a

study that level of monthly income was strongly associated with adherence to healthy diet and regular exercise³². Other studies also found that financial problem has a negative relationship with adherence to dietary^{41,42} and medical recommendations⁴³.

4. Education: Effect of education on lifestyle is as great as level of income. Education also helps support and sustains healthy lifestyle and healthy choices, fostering human development, human relationships and personal, family and community well-being. Several studies indicated that educational level of individuals was shown to be positively related with adherence to dietary regimen^{30,44}, physical activity^{28,32,44}, medications^{22,32}, smoking cessation^{22,32} and overall lifestyle recommendations⁴⁴.

II. Cognitive factors

1. Beliefs: Beliefs such as treatment can have some benefits⁴⁵, a disease can have serious consequences^{46,47}, they are susceptible to a disease⁴⁸, they have the ability to carry out the prescribed treatment regimen^{49,50} was positively associated with adherence to overall lifestyle changes. Negative health belief had negative association with dietary adherence^{41,42}. Patients' health beliefs are affected by their health literacy, and these beliefs are also contributors to non-adherence to medications⁵¹. If patients hold beliefs that are incongruent with their physician's prescription, or if their family or colleagues hold contradictory views about their illnesses and treatments, patients may have trouble even forming a willingness or intention to adhere^{52,53}.

2. Perceptions: Perceptions go on to shape a person's behavioral intentions and, ultimately, their subsequent actions. It was revealed in studies that negative perceptions were associated with non-adherence to dietary regimen^{41,42}. A perception that exercise exacerbated their illness, found negatively associated with adherence to physical activity⁵⁴.

3. Knowledge: Knowledge about healthy food, nutrition, benefits of exercise, and harmful effects of tobacco and alcohol

consumption, consequences of non-adherence to healthy lifestyle is helpful in increasing adherence of the people to healthy lifestyle. Knowledge has Positive relationship with adherence to diet^{31,55}, physical activity⁵⁵, medical adherence^{43,55}, and overall lifestyle changes^{55,56}.

4. Awareness about health benefits of healthy lifestyle and medications: A study on diabetics found positive relationship between awareness about benefits of medications and medical adherence²². Lack of information was an important reason of non-adherence to diet regimen^{41,42,54} and physical activity⁵⁴. Sufficient information provided by doctors on benefits and use of prescribed medicine and proposed lifestyle changes helps in adherence to lifestyle recommendations^{35,56}.

5. Motivation: Motivation found to have positive relationship with overall lifestyle changes^{57,58,59}. Low outcome expectancies serve as demotivating factors. It was found in a study that lack of motivation was associated with non-adherence to exercise^{60,61}. Exercising with partner can enhance motivation and support. Exercising with a partner can be more fun and competition make people perform better. Numerous studies revealed that lack of an exercise partner is associated with non-adherence to physical activity recommendations^{2,41,54,60}.

6. Consciousness and self-efficacy: Consciousness and self-efficacy found to have positive relationship with adherence to lifestyle changes *i.e.* healthy diet and physical exercise⁶². A sense of self-efficacy found to strengthen motivation to adhere to healthy lifestyle⁶³. Self-efficacy for physical activity was found as one of the facilitators of physical exercise⁶⁴.

7. Poor memory: An important factor influencing adherence was patients' ability to remember the details of the recommendations made to them. Studies have repeatedly shown that forgetting to take medications was a major contributor to non-adherence^{65,66,67}.

III. Interpersonal factors

1. Social support: The social environment and the social support available to patients also

affect their willingness to adhere to recommended treatment regimen^{68,69}. Studies have found that social support was positively related with adherence to dietary and physical activity recommendations^{70,71,72} and overall lifestyle changes^{73,74} but there is a study which found no significant relationship found between social support and overall lifestyle changes⁵⁰.

2. Lack of family support: It was found that lack of support from spouse and family was associated with non-adherence to dietary recommendation^{41,42,75}. A study says that family conflicts were associated with non-adherence to lifestyle recommendations⁷⁶. Family support was found to be the strongest and most consistent predictor of adherence to treatment in patients with type 2 diabetes⁷⁷.

3. Relationship between health-care providers and patient: A good relationship between health-care provider and patient was found to be associated with adherence to lifestyle recommendations^{56,78}. Patients reporting high levels of concordance with the physician were more likely to be compliant in taking medications prescribed during consultation⁷⁹. Patient's involvement in decision making helps in increasing patient's adherence to medications and lifestyle change recommendations^{21,80}.

4. Trust in physician: Patients' trust in their physicians is essential to their emotional disclosure and is therefore a crucial component of the patient–physician relationship. Patients must believe that their physician is someone who can understand their unique experience of being a patient, and someone who can provide them with reliable and honest advice. Trusting relationships between physicians and patients can greatly affect patient outcomes. It has been shown that physicians who promote trust in the therapeutic relationship, and who express compassion for their patients succeeds in fostering patient adherence with a variety of preventive and treatment recommendations. Various studies revealed that trust in physician was positive related with adherence to lifestyle changes^{35,78,79}.

5. Patient-physician communication: Patients who feel that their physicians communicate well with them and actively encourage them to be involved in their own care tend to be more motivated to adhere^{82,83}. Additionally, when physicians and patients agree on how involved patients should be in their care, adherence is improved. Effective interpersonal communication makes it possible for patients and physicians to work together to help patients follow mutually agreed-upon recommendations, promotes greater patient satisfaction which in turn fosters higher levels of adherence⁸⁴. Poor communication between patient-health care practitioners is associated with poorer treatment adherence in patients with diabetes¹⁵.

IV. Unintentional/ other factors

1. Busy schedule: Numerous studies revealed that busy schedule was associated with non-adherence to physical activity recommendations. Work responsibilities, family responsibilities, obligations and lack of time makes it difficult to adhere to regular physical exercise^{2,41,60,85,86,87,88}.

2. Health: Studies indicated that better health was associated with adherence to physical activity recommendations^{90,91} and poor health was associated with non-adherence to physical activity^{2,41,88,92} and overall lifestyle recommendations⁴⁶. But few researchers also found that those who perceive themselves to be in poor health were less likely than those with good health to adhere to lifestyle changes^{32,35}.

5. Unwillingness: An important reason for non-adherence to lifestyle changes that individuals do not consider the need for change, are resistant to suggestions of change, and being not willing to invest the necessary effort to achieve the desired outcome. Unwillingness is associated with non-adherence to dietary^{41,42,88} and physical activity recommendations^{2,41}.

6. Smoking: Smoking was found to be associated with non-adherence to physical activity⁸⁹. In a study conducted in Finland in hypertension patients, non-smokers were more compliant to the diet restrictions⁹³. Several Studies about compliance indicated that patients who smoked or drank alcohol were more likely to be non-compliant to the medications or treatment^{94,95,96,97}.

7. Extreme weather conditions (hot/cold/rainy season): The weather in which people exercise is very influential in terms of exercise adherence. It can often be the case that inclement weather can prevent people from exercising. Exercising in inclement weather may also compromise health. Poor weather was found to have negative relationship with adherence to physical activity recommendation^{2,41}. Intensely hot summer weather was found to be associated with non-adherence to exercise⁸⁸. Patients reported in a study that surrounding environment or weather doesn't suit them to exercise regularly⁸⁵.

8. Side effect: Side effect of medicines found to have a negative relationship with adherence to medical recommendations^{22,98}. Numerous studies on side effects factor found that side effects threaten patient's compliance to medications^{99,100,101,102}.

9. Cost of the treatment: Cost of prescribed drugs has negative relationship with adherence to medical recommendations^{35,103}. For many patients, if income is low, cost of medication directly affects their level of medication adherence especially when it is a long term therapy^{19,104,105}.

10. Feeling better/worse: Feeling better or worse was found to be associated with non-adherence to medical recommendations. If patient feel better or worse after taking medications, in both conditions patient found to become non-compliant to medications⁴³. A research found significant positive association between sense of normality and motivation to adhere to healthy lifestyle⁵⁷.

11. Eating out: Increasing number of fast food outlet and high frequency of consumption of fast food found to be a reason for non-

adherence to diet regimen^{42,88}. People who have tendency to eat out adhere less to dietary recommendations⁵⁴. Patients reported in a study that situational factors like eating out at restaurant and inappropriate foods offers by others affect their recommended diet plan⁸⁵.

12. Social gatherings: Social get-togethers are frequent in prosperous communities where energy rich and fatty food items are served. Social gatherings found to be associated with non-adherence to healthy diet⁸⁸. Most of the patients reported in a study that high frequency of social gathering (functions or festivals) with family and friends affect their healthy diet plan⁸⁵.

13. Depression: Depression has long been known to predict poor health outcomes, a fact that may be explained partly by the adherence problems caused by depression. Depressed patients experience pessimism, cognitive impairments, and withdrawal from social support, all of which can diminish both the willingness and ability to follow treatment regimens. A research suggests that one of the strongest predictors of patient non-adherence to medical treatment is patients' depression¹⁰⁶.

Studies indicated that depressed people were more likely to be non-adherent to weight loss diet regimen⁶⁰ and overall lifestyle changes^{11,107}.

14. Stress: Various stressors such as long working hours, shift work, responsibilities and a continually changing work environment may affect adoption of healthy lifestyle behaviors¹⁰⁸. Researches revealed that stress was one of the reason for non-adherence to weight loss diet⁶⁰ and overall lifestyle changes⁸⁸.

Ways to improve adherence to lifestyle changes

Health professionals can improve adherence by tailoring their communications according to the individual patient's knowledge, understanding, beliefs, preferences, needs and circumstances; and maintaining motivation by emphasizing the role of therapy in reducing cardiovascular risk. Effective intervention need not be excessively time consuming. One should ensure that cost-effective drugs are

prescribed to the patients²². Decreasing number of medicine doses or using simple reminder system may be helpful for some patients, arranging follow-up test or consultation may assist with adherence to medicines¹⁰⁹. Adherence can be improved by patient education, motivational strategies, and improving doctor-patient relationship¹¹⁰.

CONCLUSIONS

Level of non-adherence is high whether it is lifestyle recommendation or treatment prescription. Numerous factors are responsible for non-adherence, and for each type of recommendation, factors are different. A health professional should consider all possible factors which hinders adherence of the patients to the diet, physical activity, and treatment regimen while guiding the patients. There is a need for education to address the lack of information on benefits of treatment and healthy diet as well as the benefits of exercise and how exercise should be undertaken.

REFERENCES

1. Burnet, D.L., Elliott, L.D., Quinn, M.T., Plaut, A.J., Schwartz, M.A., Chin, M.H., Preventing diabetes in the clinical setting, *J Gen Intern Med.* **21**: 84–93 (2006).
2. Wadden, T.A., West, S.D., Delahanty, L. Jakicic, J., Rejeski, J., Williamson, D., Berkowitz, R.I., Kelley, D.E., Tomchee, C., Hill, J.O., Kumanyika, S. The Look AHEAD Study: A descriptive of the lifestyle intervention and the evidence supporting it, *Obesity (Silver Spring)* **14**(5): 737–752 (2006).
3. Bazzano, L.A., Serdula, M., Liu, S., Prevention of type 2 diabetes by diet and lifestyle modification, *J Am Coll Nutr.* **24**: 310–319(2005).
4. Harris, S.B., Petrella, R.J., Leadbetter, W., Lifestyle interventions for type 2 diabetes-relevance for clinical practice, *Can Fam Physician* **49**: 1618–1625 (2003).
5. The Diabetes Prevention Program (DPP) Research Group, The Diabetes Prevention Program (DPP): Description of lifestyle intervention, *Diabetes Care* **25**: 2165–2171 (2002).
6. Wadden, T.A., Webb, V.L., Moran, C.H., Bailer, B.A., Lifestyle Modification for Obesity: New Developments in Diet, Physical Activity, and Behavior Therapy, *Circulation* **125**(9): 1157-1170 (2012).
7. Foster-Schubert, K.E., Alfano, C.M., Duggan, C.R., Xiao, L., Campbell, K.L., Kong, A., Bain, C., Wang, C.Y., Blackburn, G., and McTiernan, A., Effect of diet and exercise, alone or combined, on weight and body composition in overweight-to-obese postmenopausal women, *Obesity (Silver Spring)* **20**(8): 1628–1638 (2012).
8. Lehane, E., McCarthy, G., Intentional and unintentional medication non-adherence: A comprehensive framework for clinical research and practice? A discussion paper, *Int J Nurs Stud.* **44**(8): 1468-1477 (2007).
9. DiMatteo, M.R., Variations in patients' adherence to medical recommendations: A quantitative review of 50 years of research, *Medical Care* **42**: 200-209 (2004a).
10. DiMatteo, M.R., Reiter, R.C., and Gambone, J.C., Enhancing medication adherence through communication and informed collaborative choice, *Health Communication* **6**: 253-265. (1994).
11. Haynes, R.B., McDonald, H.P., & Garg, A.X., Helping patients follow prescribed treatment: Clinical applications, *Journal of the American Medical Association* **288**: 2868-2879 (2002).
12. Park, K., Cho, S., Bower, J.K., Changes in Adherence to Non-Pharmacological Guidelines for Hypertension, *PLoS One* **11**(8): e0161712 (2016).
13. Hernandez-Ronquillo, L., Tellez-Zenteno, J.E., Garduno-Espinosa, J., González-Acevez, E., Factors associated with therapy noncompliance in type 2 diabetes patients, *Salud Publica de Mexico* **45**(3): 191-197 (2003).
14. Melikian, C., White, T.J., Vanderplas, A., Dezii, C.M., Chang, E., Adherence to oral diabetic therapy in a managed care

- organization, *Clin Ther.* **24(3)**: 460-467 (2002).
15. Ciechanowski, P.S., Katon, W.J., Russo, J.E., Walker, E.A., The patient-provider relationship: attachment theory and adherence to treatment in diabetes, *American Journal of Psychiatry* **158(1)**: 29-35 (2001).
 16. Ho, P.M., Magid, D.J., Shetterly, S.M., Olson, K.L., Peterson, P.N., Masoudi, F.A., Rumsfeld, J.S., Importance of therapy intensification and medication nonadherence for blood pressure control in patients with coronary disease, *Arch Intern Med.* **168**: 271–276 (2008).
 17. Nelson, M.R., Reid, C.M., Ryan, P., Willson, K., Yelland, L., Self-reported adherence with medication and cardiovascular disease outcomes in the Second Australian National Blood Pressure Study (ANBP2), *Med J Aust.* **185(9)**: 487-489 (2006).
 18. Cutrona, S.L., Choudhry, N.K., Stedman, M., Servi, A., Liberman, J.N., Brennan, T., Shrank, W. H., Physician effectiveness in interventions to improve cardiovascular medication adherence: A systematic review, *Journal of General Internal Medicine* **25**: 1090-1096 (2010).
 19. World Health Organization, Adherence to long-term therapies: evidence for action, Geneva, (2003). <http://apps.who.int/iris/bitstream/10665/42682/1/9241545992.pdf>
 20. Bondesson, A., Hellström, L., Eriksson, T., and Höglund, P., A structured questionnaire to assess patient compliance and beliefs about medicines taking into account the ordered categorical structure of data, *Journal of Evaluation in Clinical Practice* **15**: 713-723 (2009).
 21. Arbuthnott, A., and Sharpe, D., The effect of physician-patient collaboration on patient adherence in non-psychiatric medicine. *Patient Education and Counselling* **77**: 60-67 (2009).
 22. Sharma, T., Kalra, J., Dhasmana, D.C., Basera, H., Poor adherence to treatment: A major challenge in diabetes, *JACM* **15(1)**: 26-29 (2014).
 23. De Geest, S., Von Renteln-Kruse, W.V., Steeman, E., Degraeve, S., and Abraham, I.L., Compliance issues with the geriatric population, *Nursing Clinics of North America* **33**: 467-480 (1998).
 24. Howarth, N.C., Huang, T.T., Roberts, S.B., Lin, B.H., McCrory, M.A., Eating patterns and dietary composition in relation to BMI in younger and older adults, *Int J Obes (Lond)* **31**:675– 684 (2007).
 25. Wu, S.J., Chang, Y.H., Wei, I.L., Kao, M.D., Lin, Y.C., Pan, W.H. Intake levels and major food sources of energy and nutrients in the Taiwanese elderly, *Asia Pac J Clin Nutr.* **14(3)**: 211-220 (2005).
 26. Stables, G.J., Subar, A.F., Patterson, B.H., Dodd, K., Heimendinger, J., Van Duyn, M.A., Nebeling, L., Changes in vegetable and fruit consumption and awareness among US adults: results of the 1991 and 1997 5 A Day for Better Health Program surveys, *J Am Diet Assoc.* **102(6)**: 809-817 (2002).
 27. Steptoe, A., Doherty, S., Kerry, S., Rink, E., Hilton, S., Sociodemographic and psychological predictors of changes in dietary fat consumption in adults with high blood cholesterol following counseling in primary care, *Health Psychol.* **19**: 411–419 (2000).
 28. Uzun, S., Kara, B., Yokuşoğlu, M., Arslan, F., Yilmaz, M.B., Karaeren, H., The assessment of adherence of hypertensive individuals to treatment and lifestyle change recommendations, *Anadolu Kardiyol Derg.* **9(2)**: 102-109 (2009).
 29. Horne, R., and Weinman, J., Patients' beliefs about prescribed medicines and their role in adherence to treatment in chronic physical illness, *Journal of Psychosomatic Research* **47**:555-567 (1999).
 30. Mumu, S.J., Saleh, F., Ara, F., Afnan, F., Ali, L., Non-adherence to life-style modification and its factors among type 2

- diabetic patients, *Indian J. Public Health* **58**: 40-44 (2014).
31. Parajuli, J., Saleh, F., Thapa, N., Ali, L., Factors associated with non-adherence to diet and physical activity among Nepalese type 2 diabetes patients; a cross sectional study, *BMC Res Notes* **7**:1-9 (2014).
32. Elbur, A.I., Level of Adherence to Lifestyle Changes and Medications among Male Hypertensive Patients in Two Hospitals in Taif; Kingdom Of Saudi Arabia, *International Journal of Pharmacy and Pharmaceutical Sciences* **7(4)**:168-172 (2015).
33. Aggarwal, B., Mosca, L., Lifestyle and psychosocial risk factors predict non-adherence to medication, *Annals of Behavioral Medicine* **40**: 228-233 (2010).
34. Jerant, A., Chapman, B., Duberstein, P., Robbins, J., & Franks, P., Personality and medication non-adherence among older adults enrolled in a six-year trial, *British Journal of Health Psychology* **16**: 51-169 (2010).
35. Levesque, A., Li, H.Z., Pahal, J.S., Factors Related to Patients' Adherence to Medication and Lifestyle Change Recommendations: Data from Canada, *International Journal of Psychological Studies* **4(2)**: 42-55 (2012).
36. Forshee, R.A., Storey, M.L., Demographics, not beverage consumption, is associated with diet quality, *Int J Food Sci Nutr.* **57**: 494 -511 (2006).
37. Hart, A. Jr., Tinker, L., Bowen, D.J., Longton, G., Beresford, S.A., Correlates of fat intake behaviors in participants in the eating for a healthy life study, *J Am Diet Assoc.* **106**: 1605–1613 (2006).
38. Nasreddine, L., Hwalla, N., Sibai, A., Hamze, M., Parent-Massin, D., Food consumption patterns in an adult urban population in Beirut, Lebanon, *Public Health Nutr.* **9(2)**: 194-203 (2006).
39. Satia, J.A., Galanko, J.A., Neuhaus, M.L., Food nutrition label use is associated with demographic, behavioral, and psychosocial factors and dietary intake among African Americans in North Carolina, *J Am Diet Assoc.* **105**: 392–402(2005).
40. Dynesen, A.W., Haraldsdóttir, J., Holm, L., Astrup, A., Sociodemographic differences in dietary habits described by food frequency questions: results from Denmark, *Eur J Clin Nutr.* **57**: 1586 – 1597 (2003).
41. Khan, A.R., Al-Abdul Lateef, Z.N., Al Aithan, M.A., Bu-Khamseen M.A., Al Ibrahim, I., Khan, S.A., Factors contributing to non-compliance among diabetics attending primary health centers in the Al Hasa district of Saudi Arabia, *J Family Community Med.* **19(1)**: 26-32(2012).
42. Misra, A., Khurana, L., Obesity and the metabolic syndrome in developing countries, *J Clin Endocrinol Metab.* **93 (11 Suppl 1)**: S9-S30 (2008).
43. Sajith, M., Pankaj, M., Pawar, A., Modi, A., Sumariya, R., Medication Adherence to Antidiabetic Therapy in Patients with Type 2 Diabetes Mellitus, *Int J Pharm Pharm Sci.* **6(Suppl 2)**: 564-570 (2014).
44. Sováriová Soósová, M., Hrehová, J., The Effect of Education on Lifestyle Changes and Metabolic Syndrome Components, *Cent Eur J Nurs Midw.* **5(4)**: 161-168 (2014).
45. Foster, J.M., Smith, L., Bosnic-Anticevich, S.Z., Usherwood, T., Sawyer, S.M., Rand, C.S., and Reddel, H.K. Identifying patient-specific beliefs and behaviours for conversations about adherence in asthma. *Intern Med J.* **42(6)**: e136-144 (2011).
46. DiMatteo, M.R., Haskard, K.B., and Williams, S.L., Health beliefs, disease severity, and patient adherence: A meta-analysis, *Medical Care* **45**: 521-528 (2007).
47. Stafford, L., Jackson, H.J., and Berk, M., Illness beliefs about heart disease and adherence to secondary prevention regimens, *Psychosom Med.* **70**: 942-948 (2008).
48. Becker, M.H., The Health Belief Model and prediction of dietary compliance: A

- field experiment, *Journal of Health and Social Behavior* **18**: 348-366 (1977).
49. Dilorio, C., McCarthy, F., Depadilla, L., Resnicow, K., Holstad, M.M., Yeager, K., Lundberg, B., Adherence to antiretroviral medication regimens: A test of a psychosocial model, *Aids and Behavior* **13**: 10-22 (2009).
50. Brus, H., Van de Laar, M., Taal, E., Rasker, J., and Wiegman, O., Determinants of compliance with medication in patients with rheumatoid arthritis: The importance of self-efficacy expectancies, *Patient Education Counselling* **36**: 57-64 (1999).
51. Anarella, J., Roohan, P., Balistreri, E., Gesten, F., A survey of Medicaid recipients with asthma: perceptions of self-management, access, and care, *Chest* **125(4)**: 1359–1367 (2004).
52. Soliday, E., Hoeksel, R., Health beliefs and pediatric emergency department after-care adherence, *Ann Behav Med.* **22(4)**: 299-306 (2000).
53. Straughan, P.T., Seow A., Attitudes as barriers in breast screening: a prospective study among Singapore women, *Soc Sci Med.* **51(11)**: 1695–1703 (2000).
54. Ganiyu, A.B., Mabuza, L.H., Malete, N.H., Govender, I., Ogunbanjo, G.A. Non-adherence to diet and exercise recommendations amongst patients with type 2 diabetes mellitus attending Extension II Clinic in Botswana, *African Journal of Primary Health Care & Family Medicine*, **5(1)**: 1-6 (2013).
55. Alm-Roijer, C., Stagmo, M., Udén, G., Erhardt, L., Better Knowledge Improves Adherence to Lifestyle Changes and Medication in Patients with Coronary Heart Disease, *Eur J Cardiovasc Nurs.* **3(4)**: 321-330 (2004).
56. Gellad, W.F., Grenard, J.L., and Marcum, Z.A., A systematic review of barriers to medication adherence in elderly: Looking beyond cost and regimen complexity, *The American Journal of Geriatric Pharmacotherapy* **9**: 11-23 (2011).
57. Kahkonen, O., Kankkunen, P., Saaranen, T., Miettinen, H., Kyngas, H. and Lamidi, M.L. Motivation Is A Crucial Factor For Adherence To A Healthy Lifestyle Among People With Coronary Heart Disease After Percutaneous Coronary Intervention. *J Adv Nurs.* **71(10)**: 2364-2673 (2015).
58. Zarani, F., Sarami, G., Sadeghian, S., Adherence in CABG Patients: An Emperical Test of Health Behaviour Model. *International Journal of clinical Medicine* **5**: 225-233 (2014).
59. Kaariainen, M., Paukama, M., and Kyngas, H., Adherence with health regimens of patients on Warfarin therapy, *Journal of Clinical Nursing* **22**: 89–96 (2013).
60. Sharifi, N., Mahdavi, R., and Ebrahimi-Mameghani, M., Perceived Barriers to Weight loss Programs for Overweight or Obese Women, *Health Promot Perspect.* **3(1)**: 11-22 (2013).
61. Beverly, E.A., Wray, L.A., The role of collective efficacy in exercise adherence: a qualitative study of spousal support and Type 2 diabetes management, *Health Educ. Res.* **25(2)**: 211-223 (2010).
62. Martos-Méndez, M.J., Self-Efficacy and Adherence to Treatment: The Mediating Effects of Social Support, *Journal of Behavior, Health & Social Issues* **7(2)**: 19-29 (2015).
63. Artinian, N.T., Fletcher, G.F., Mozaffarian, D., Kris-Etherton, P., Van Horn, L., Lichtenstein, A.H., Kumanyika, S., Kraus, W.E., Fleg, J.L., Redeker, N.S., Meininger, J.C., Banks, J., Stuart-Shor, E.M., Fletcher, B.J., Miler, T.D., Hughes, S., Braun, L.T., Kopin, L.A., Berra, K., Hayman, L.L., Ewing, L.J., Ades, P.A., Durstine, J.L., Houston-Miller, N. and Burke, L.E., Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a scientific statement from the American Heart Association, *Circulation* **122**: 406–441 (2010).
64. Stutts, W.C., Physical activity determinants in adults. Perceived benefits,

- barriers, and self efficacy, *AAOHN J.* **50(11)**: 499-507 (2002).
65. Brekke, H.K., Sunesson, A., Axelsen, M., Lenner, R.A., Attitudes and barriers to dietary advice aimed at reducing risk of type 2 diabetes in first-degree relatives of patients with type 2 diabetes, *J Hum Nutr Diet* **17**: 513–521 (2004).
66. Shemesh, E., Shneider, B.L., Savitzky, J.K., Arnott, L., Gondolesi, G.E., Krieger, N.R., Kerkar, N., Magid, M.S., Stuber, M.L., Schmeidler, J., Yehuda, R., Emre, S., Medication adherence in pediatric and adolescent liver transplant recipients, *Pediatrics* **113(4)**: 825–832 (2004).
67. Zaghloul, S.S., Goodfield, M.J., Objective assessment of compliance with psoriasis treatment, *Arch Dermatol*, **140(4)**: 408-414 (2004).
68. Kadam, U.T., Croft, P., McLeod, J., Hutchinson, M., A qualitative study of patients' views on anxiety and depression, *Br J Gen Pract.* **51**: 375–380 (2001).
69. Sirey, J.A., Bruce, M.L., Alexopoulos, G.S., Perlick, D.A., Friedman, S.J., Meyers, B.S., Stigma as a barrier to recovery: perceived stigma and patient-rated severity of illness as predictors of antidepressant drug adherence, *Psychiatr Serv.* **52(12)**: 1615–1620 (2001).
70. Engbers, L.H., van Poppel, M.N., Chin, A. Paw, M., van Mechelen, W., The effects of a controlled worksite environmental intervention on determinants of dietary behavior and self-reported fruit, vegetable and fat intake, *BMC Public Health* **6**: 253 (2006).
71. Kahn, E.B., Ramsey, L.T., Brownson, R.C., Heath, G.W., Howze, E.H., Powell, K.E., Stone, E.J., Rajab, M.W., Corso, P., The effectiveness of interventions to increase physical activity: a systematic review, *Am J Prev Med.* **22**: 73–107(2002).
72. Trost, S.G., Owen, N., Bauman, A.E., Sallis, J.F., Brown W., Correlates of adults' participation in physical activity: review and update, *Med Sci Sports Exerc.* **34(12)**: 1996 -2001(2002).
73. Karamanidou, C., Clatworthy, J., Weinman, J., and Horne, R., A systematic review of the prevalence and determinants of nonadherence to phosphate binding medication in patients with end-stage renal disease, *BMC Nephology* **9**: 2 (2008).
74. Molloy, G.J., Perkins-Porras, L., Bhattacharyya, M.R., Strike, P.C., and Steptoe, A., Practical support predicts medication adherence and attendance at cardiac rehabilitation following acute coronary syndrome, *J Psychosom Res.* **65(6)**: 581-586 (2008).
75. Nicklett, E., Liang, J., Diabetes-related support, regimen adherence, and health decline among older adults, *J Gerontol B Psychol Sci Soc Sci.* **65B(3)**: 390–399 (2010).
76. DiMatteo, M.R., Social support and patient adherence to medical treatment: A meta-analysis. *Health Psychology* **23**: 207-218 (2004b).
77. Glasgow, R.E., Toobert, D.J., Social environment and regimen adherence among type II diabetes patients, *Diabetes Care* **11**: 377–386 (1988).
78. Bennet, J.K., Fuertes, J.N., and Phillips, R., The role of patient attachment and working alliance on patient adherence, satisfaction, and health-related quality of life in lupus treatment, *Patient Education and Counseling* **85**: 53-59 (2010).
79. Kerse, N., Buetow, S., Mainous, A.G. 3rd, Young, G., Coster, G., and Arroll, B., Physician-patient relationship and medication compliance: A primary care investigation, *Ann Fam Med.* **2(5)**: 455-461 (2004).
80. Bultman, D.C., Svarstad, B.L., Effects of physician communication style on client medication beliefs and adherence with antidepressant treatment, *Patient Education and Counseling*, **40**: 173-185 (2000).
81. Branch, W.T. Jr., The ethics of caring and medical education, *Acad Med.* **75(2)**: 127–132 (2000).
82. O'Malley, A.S., Forrest, C.B., Mandelblatt, J., Adherence of low-income

- women to cancer screening recommendations, *J Gen Intern Med.* **17(2)**: 144-154 (2002).
83. Martin, L.R., DiMatteo, M.R., Lepper, H.S., Facilitation of patient involvement in care: development and validation of a scale, *Behav Med.* **27**: 111–120 (2001).
84. Jahng, K.H., Martin, L.R., Golin, C.E., DiMatteo, M.R., Preferences for medical collaboration: patient-physician congruence and patient outcomes, *Patient Educ Couns.* **57(3)**: 308–314 (2005).
85. Jadawala, H.D., Pawar, A.B., Patel, P.B., Patel, K.G., Patel, S.B., Bansal, R.K., Factors Associated With Non Adherence to Diet and Physical Activity among Diabetes Patients: A Cross Sectional Study, *National Journal of Community Medicine* **8(2)**: 68-73 (2017).
86. Thomas, N., Alder, E., Leese, G.P., Barriers to physical activity in patients with diabetes, *Postgrad Med J.* **80(943)**: 287–291 (2004).
87. Wanko, N.S., Brazier, C.W., Young-Rogers, D., Dunbar, V.G., Boyd, B., George, C.D., Rhee, M.K., el-Kebbi, I.M., Cook, C.B. Exercise preferences and barriers in urban African Americans with type 2 diabetes. *Diabetes Educ.* **30(3)**: 502-513 (2004).
88. Serour, M., Alqhenaei, H., Al-Saqabi, S., Mustafa, A., Ben-Nakhi, A., Cultural factors and patients' adherence to lifestyle measures, *Br J Gen Pract.* **57(537)**: 291–295 (2007).
89. Ahmed, A.T., Karter, A.J., Liu, J., Alcohol consumption is inversely associated with adherence to diabetes self-care behaviours, *Diabet Med.* **23(7)**: 795–802 (2006).
90. Ainsworth, B.E., Wilcox, S., Thompson, W.W., Richter, D.L., Henderson, K.A., Personal, social, and physical environmental correlates of physical activity in African-American women in South Carolina, *Am J Prev Med.* **25(suppl 1)**: 23–29 (2003).
91. Norman, A., Bellocco, R., Vaida, F., Wolk, A., Total physical activity in relation to age, body mass, health and other factors in a cohort of Swedish men, *Int J Obes* **26(5)**: 670-675 (2002).
92. Booth, M.L., Owen, N., Bauman, A., Clavisi, O., Leslie, E., Social-cognitive and perceived environment influences associated with physical activity in older Australians, *Prev. Med.* **31**: 15–22 (2000).
93. Kyngas, H., Lahdenpera, T., Compliance of patients with hypertension and associated factors, *J Ad Nurs.* **29(4)**: 832–839 (1999).
94. Moreno, J.L. Catley, D., Lee, H.S., Goggin, K., The relationship between ART adherence and smoking status among HIV+ individuals, *AIDS Behav.* **19(4)**: 619-625 (2015).
95. Balbay, O., Annakkaya, A.N., Arbak, P., Bilgin, C., Erbas, M., Which patients are able to adhere to tuberculosis treatment? A study in a rural area in the northwest part of Turkey, *Jpn J Infect Dis.* **58**: 152–158 (2005).
96. Cooper, C., Carpenter, I., Katona, C., Schroll, M., Wagner, C., Fialova, D., Livingston, G., The AdHOC study of older adults' adherence to medication in 11 countries, *Am J Geriatr Psychiatry* **13**: 1067–1076 (2005).
97. Fodor, G.J., Kotrec, M., Bacskai, K., Dorner, T., Lietava J., Is interview a reliable method to verify the compliance with antihypertensive therapy? An international central-European study, *J Hypertens.* **23**: 1261–1266 (2005).
98. Kassahun, A., Gashe, F., Mulisa, E., Rike, W.A., Nonadherence and factors affecting adherence of diabetic patients to anti-diabetic medication in Assela General Hospital, Oromia Region, Ethiopia, *Journal of Pharmacy & Bioallied Sciences* **8(2)**: 124-129 (2016).
99. Lihara, N., Tsukamoto, T., Morita, S., Miyoshi, C., Takabatake, K., Kurosaki, Y., Beliefs of chronically ill Japanese patients that lead to intentional non-adherence to medication, *J Clin Pharm Ther.* **29**: 417–424 (2004).
100. Kaplan, R.C., Bhalodkar, N.C., Brown, E.J. Jr., White, J., Brown, D.L., Race,

- ethnicity, and sociocultural characteristics predict noncompliance with lipid-lowering medications, *Prev Med.* **39**:1249–1255 (2004).
101. O'Donoghue, M.N., Compliance with antibiotics, *Cutis* **73(Suppl 5)**: 30–32 (2004).
102. Ponnusankar, S., Surulivelrajan, M., Anandamoorthy, N., Suresh, B., Assessment of impact of medication counseling on patients' medication knowledge and compliance in an outpatient clinic in South India, *Patient Educ Couns.* **54(1)**:55-60 (2004).
103. Ohane Buabang, K., Motowe, L., Plange-Rhule, J., Unaffordable drug prices: the major cause of non-compliance with hypertensive medication in Ghana, *J Pharm Pharm Sci.* **7(3)**: 350-352 (2004).
104. Briesacher, B.A., Gurwitz, J.H., Soumerai, S.B., Patients at-risk for cost-related medication nonadherence: a review of the literature, *J Gen Intern Med.* **22(6)**: 864-871 (2007).
105. Piette, J.D., Wagner, T.H., Potter, M.B., Schillinger, D., Health insurance status, cost-related medication underuse, and outcomes among diabetes patients in three systems of care, *Med Care* **42(2)**: 102-109 (2004).
106. DiMatteo, M.R., Lepper, H.S., Croghan, T.W., Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence, *Arch Intern Med.* **160(14)**: 2101-2107 (2000).
107. Rejeski, W.J., Miller, M.E., King, A.C., Studenski, S.A., Katula, J.A., Fielding, R.A., Glynn, N.W., Walkup, M.P., Ashmore, J.A., Predictors of adherence to physical activity in the Lifestyle Interventions and Independence for Elders pilot study (LIFE-P), *Clin Interv Aging* **2(3)**: 485-494 (2007).
108. Han, K., Tinkoff, A.M., Storr, C.L., and Geiger-Brown, J., Job stress and work schedules in relation to nurse obesity, *Journal of Nursing Administration* **41(11)**: 488-495 (2011).
109. Benner, J.S., Tierce, J.C., Ballantyne, C.M., Prasad, C., Bullano, M.F., Willey, V.J., Follow up Lipid tests and Physician Visits Associated with Improved Adherence Statin Therapy, *Pharmacoeconomics* **22 (suppl 3)**: 13–23 (2004).
110. World Health Organization: Diabetes Mellitus: report of a WHO study group, Geneva, Switzerland: World Health Organization, Technical Report Series No. 727 (1985).