

## Effects of Seed Pelleting, Polymer Coating and Packaging Materials on Seed Quality Characters of Tomato (*Lycopersicon esculentum* L.) Seeds after Three Months of Ambient Storage

Dheeraj, M. \*, Dayal, A., Neha Thomas and Ramteke, P. W.

Dept of Genetics and Plant Breeding

Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad (U.P) - 211007

\*Corresponding Author E-mail: [dheerajtony77@gmail.com](mailto:dheerajtony77@gmail.com)

Received: 5.03.2018 | Revised: 14.04.2018 | Accepted: 22.04.2018

### ABSTRACT

The present storage experiment was conducted at Department of Genetic and Plant Breeding, Sam Higginbottom University of Agriculture, Technology and Sciences, Uttar Pradesh during 2017–2018 with tomato seeds (Pusa Ruby). The seeds were coated with polymer in combination with fungicide and maintained untreated seeds (control) where  $T_0$  is control,  $T_1$  Acacia (Acacia nilotica) leaf powder@4mg/kg seed,  $T_2$ -Imidacloprid @ 3ml/kg seed,  $T_3$ -Neem leaf powder @ 10 g per kg of seed,  $T_4$  ZnSO<sub>4</sub>@ 3.0mg/kg seed,  $T_5$  Polymer @ 2ml/kg seed, Acacia leaf powder+ Polymer @ 4ml/kg seed and  $T_7$ -Imidacloprid +polymer@ 4ml/kg seed,  $T_8$ -Imidacloprid +polymer@6ml/kg seed,  $T_9$ -Trichoderma viridae @4gm/kg seed  $T_{10}$ -Neem leaf powder+polymer@8ml/kg seed  $T_{11}$ -Vitavax+polymer @6ml/kg seed. Data was subjected to factorial experiment laid out in completely randomized. Treated seeds design were packed in aluminium foil pouch and high density polythene bag (factor  $P_1$  and  $P_2$ ) after three months and assessment of seed quality was done in terms of germination per cent, seedling length, seedling dry weight, seedling vigour indices, speed of germination, moisture content and electrical conductivity where. Per cent germination, seedling length, seedling dry weight, seedling vigour indices showed maximum in treatment  $T_{11}$ -Vitavax+polymer @6ml/kg as compared to other treatments. However, electrical conductivity and moisture content were lowest in  $T_{11}$ -Vitavax+polymer @6ml/kg.

**Key words:** Tomato, Vitavax, Imidacloprid, Acacia leaf powder, Polymer, Neem leaf powder aluminium foil pouch, High density polythene bag,

### INTRODUCTION

Tomato (*Lycopersicon esculentum* L.)  $2n=24$  a self-pollinated crop is one of the important solanaceous vegetable crops grown widely all over the world because of its special nutritive value and also its wide spread production. It is the world's largest vegetable crops next to

potato and sweet potato, but it tops the list of commercial vegetables. Health wise tomatoes are one of the "Highest perch" because they contain an antioxidant "lycopene" and for its ability to reduce the risk of prostate cancer in men.

**Cite this article:** Dheeraj, M., Dayal, A., Thomas, N., and Ramteke, P.W., Effects of Seed Pelleting, Polymer Coating and Packaging Materials on Seed Quality Characters of Tomato (*Lycopersicon esculentum* L.) Seeds after Three Months of Ambient Storage, *Int. J. Pure App. Biosci.* 6(2): 1511-1518 (2018). doi: <http://dx.doi.org/10.18782/2320-7051.6595>

Tomato ranks third in priority on consumption basis after Potato and Onion in India but ranks second after potato in the world. India ranks second in the area as well as in production of Tomato. The major tomato growing countries are China, USA, Italy, Turkey, India and Egypt. In India major tomato producing states are Andhrapradesh, Karnataka, Telangana, Odisha, Gujarat. Tomato is one of the important vegetable crops being cultivated throughout India. Rapid deterioration of stored vegetable seeds is a serious problem which occurs at an increasing rate in uncontrolled storage environment. Seed pelleting is the process of enclosing a seed with small quantity of inert material just large enough to produce globular unit of standard size to provide small amount of nutrients to young seedlings<sup>17,21,12</sup>. A knowledge on the use of pelleted seeds in improving the productivity of the tomato crop as well as proper storage of pelleted tomato seeds in suitable containers under ambient temperature, relative humidity and at relatively low cost with minimum deterioration in quality for a period of at least one or more season will be of immense useful to the seed industry in general and farming community in

particular. The application of polymers to seed serves as an extra exterior shell in order to give the desired seed characteristics viz., quick or delayed water uptake and enhanced germination that would be beneficial for better emergence and establishment in the given condition. Packaging materials play a major role in prolonging the shelf life of a seed during storage as they separate seeds from the surrounding environment. This increases storage or shelf life by inhibiting the growth of microorganisms and improves hygiene by reducing the danger of cross contamination. Therefore the present study was undertaken to know the effect of treatment and packaging materials on seed quality

### MATERIAL AND METHODS

After imposition of seed treatments, the treated seed along with untreated seeds (control) were packed and stored under ambient conditions in Seed Testing Laboratory, Department of Genetics and Plant Breeding, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad, Uttar Pradesh for three months.

**Variety-** Pusa Ruby

**Treatments** – Twelve (12)

T<sub>0</sub> - control

T<sub>1</sub> - Acacia (*Acacia nilotica*) leaf powder@4mg/kg seed,

T<sub>2</sub> - Imidacloprid @ 3ml /kg seed,

T<sub>3</sub> -Neem leaf powder @ 10 g per kg of seed,

T<sub>4</sub> . ZnSO<sub>4</sub>@ 3.0mg/kg seed,

T<sub>5</sub> -Polymer @ 2ml/kg seed,

T<sub>6</sub> -Acacia leaf powder+ Polymer @ 4ml/kg seed

T<sub>7</sub> -Imidacloprid +polymer@ 4ml/kg seed,

T<sub>8</sub> -Imidacloprid +polymer@6ml/kg seed,

T<sub>9</sub> -*Trichoderma viridae* @4gm/kg seed

T<sub>10</sub> -Neem leaf powder+polymer@8ml/kg seed

T<sub>11</sub> -Vitavax+polymer @6ml/kg seed)

**Packaging materials**

P1-aluminium foil pouch

P2-high density polythene bag

**Moisture content:** Seeds were stored at 8 % moisture content

**Temperature:** Ambient storage conditions.

**OBSERVATIONS**

**Germination per cent (%)**: It refers to the proportion by number of seeds which have produced seedlings classified as normal under the conditions and within the period specified that is the percentage of normal seedlings.

The equation to calculate germination percent is:

$$GP = \frac{\text{No. of seeds germinated} \times 100}{\text{Total no. of seeds}}$$

**Shoot Length (cm)**:

Ten normal seedlings used for root length measurement were also used for the measurement of shoot length. The shoot length was measured from the base of the primary leaf to the base of the hypocotyls and mean shoot length was expressed in centimeters.

**Root length (cm)**:

From the germination test, ten normal seedlings were selected randomly in each treatment from all the replication on 8th day. The root length was measured from the tip of the primary root to base of hypocotyls and mean root length was expressed in centimeters.

**Seedling length (cm)**:

Seedling length is the best indicator of seed vigour. The relative length of root and shoot of seedlings would predict their subsequent growth and performance.

**Dry matter (g)**:

Dry weight were based on weights determined before and after oven drying of seedling samples at 103<sup>0</sup>C for 4hours

**Vigour index**:

The computed vigour index, which is the totality of performance, has been regarded as a good index to measure the quality of seed lots.

**Vigour index I**: Germination (%) x seedling length (cm).

**Vigour index II**: Germination (%) x seedling dry weight (cm).

**Moisture content (%)**:

Moisture content of seed was determined as per ISTA rules. Five grams of seed was weighed, ground and placed in aluminium cups. The aluminium cups along with ground seed material was dried in hot air oven

maintained at 130± 1 <sup>0</sup>C temperature for four hours.

In this method (Air oven method) the seed moisture content is determined by removing the excess moisture content from the seeds in an air oven the submitted sample meant for moisture testing must be in an intact moisture proof container from which as much as air is excluded. The determination should be made as soon as possible. (Weight of submitted sample for tomato 70gm)

The moisture content was determined on dry weight basis using the following formula.

$$\text{Moisture content (\%)} = \frac{W_2 - W_3}{W_2 - W_1} \times 100$$

Where, W<sub>1</sub> - Weight of the empty container with its cover (g)

W<sub>2</sub>- Weight of container its cover and ground seeds before drying

W<sub>3</sub>- Weight of container its cover and ground seeds after drying

**Speed of germination**:

Seeds were germinated in paper medium with four replications of 100 seed each. The number of seeds germinated was recorded daily upto the day of final count. The speed of germination was calculated by using the formula suggested by Maguire.

Speed of germination = + + ----- Where, Xn: Number of seeds germinated at nth count  
Yn: Number of days from sowing to nth count.

**Electrical conductivity** (dSm<sup>-1</sup>) Twenty-five seeds were taken randomly from each treatment in four replicates and they were surface sterilized to remove chemical residues, if any. Then they were soaked in 100 ml distilled water for 12 hours at a temperature of 25+/-1 <sup>0</sup>C. After incubation the conductivity of seed leachate was measured in a digital conductivity meter (model-DI909/DS-7007) and the EC was expressed in dSm<sup>-15</sup>.

Experiment was laid out in completely randomized design with factorial concept in four replications. The seed samples drawn at after three months months of storage and evaluated for various seed quality parameters in order to determine the suitable treatment for better storage.

## RESULTS AND DISCUSSION

Significant results were obtained due to seed pelleting with polymer coating for the seed quality parameters. Results were significantly recorded higher in seeds coated with T<sub>11</sub>-Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>- Imidacloprid +polymer@ 4ml/kg seed stored in aluminium foil pouch compared to all other treatments and the lowest germination percent, shoot length, root length, seedling length, seedling dry weight, vigour index I and II recorded in T<sub>0</sub> (control) at the end of 3 months storage (Table). Similar results were also reported by Geetharani *et al*, Kamara *et al*.<sup>11</sup>, Almeida<sup>4</sup>. At the end of 3 months of storage period, the lowest moisture content and electrical conductivity was recorded in the seeds coated with T<sub>11</sub>- Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>- Imidacloprid +polymer@ 4ml/kg seed stored in aluminium foil pouch and the highest moisture content and electrical conductivity was recorded in T<sub>0</sub> (control).The film formed around the seed act as a physical barrier, which has been reported to reduce leaching of inhibitors from the seed covering and may restrict oxygen diffusion to the embryo.

Germination percent in T<sub>11</sub>-Vitavax+polymer @6ml/kg (88.97%) recorded higher germination because of lower respiration rate and metabolic activity and inactivation of enzymes required for retention of germination for longer period Dhatt *et al.*, it is evidenced by higher germination at end of 3 months.

Higher Shoot length(7.77), root length(10.02), Seedling length(17.97), seedling dry weight(39.80) was obtained in T<sub>11</sub>-Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>- Imidacloprid +polymer@ 4ml/kg (7.64), (9.64), (17.45), (38.78) and lower in T<sub>0</sub>P<sub>1</sub> (5.71), (8.03), (13.73), (30.55) respectively. It was due to higher percentage and better germination of seedlings in seeds coated with polymer fungicide and insecticide this protects fungi invasion thereby good and better germination and subsequent higher root and shoots lengths seedling dry weight Basavaraj *et al.* in onion. Higher seedling

length and seedling dry weight is an indication of maintenance of vigour in the seeds preserved in ambient storage,

Higher seed vigour index I(1686), II (3728) was obtained in T<sub>11</sub>-Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>-Imidacloprid +polymer@ 4ml/kg (1623), (3657), and lower in T<sub>0</sub>P<sub>1</sub> (1377), (3105), respectively in polymer coating along with fungicide and insecticide it is due to more germination, root and shoot length, seedling dry weight, lesser infection by storage fungi.

Higher speed of germination (37.94) was obtained in T<sub>11</sub>-Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>-Imidacloprid +polymer@ 4ml/kg (37.28) and lower in T<sub>0</sub>P<sub>1</sub> (35.76) it was due Coating of seeds with polymer and fungicides might have protected the seed from influence of depletion factors resulting in maintenance of seed viability for a comparatively longer period. These findings are in agreement with the results obtained by Jitendra *et al.*<sup>10</sup>, Pawar *et al.*

The lower electrical conductivity (0.394) was obtained in T<sub>11</sub>- Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>- Imidacloprid +polymer@ 4ml/kg (0.447) higher in T<sub>0</sub>P<sub>2</sub> (0.537) this was mainly governed by lower cell wall permeability which indicated lower respiration rate and metabolic activity maintenance of vigour during storage. This finding is supported by Doijode 2000 in winter squash and Kumar *et al* in karanj seeds. The Lower moisture content was observed is (8.11) in T<sub>11</sub>-Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>- Imidacloprid +polymer@ 4ml/kg (8.23) higher in T<sub>0</sub>P<sub>2</sub> (8.67) is mainly because maintenance of lower moisture content 8 % during the storage period, container viz., aluminium foil pouch and high density polythene bag acted as moisture proof containers this is evidenced by lower moisture content 8% in the seeds packed in aluminium foil pouch and high density polythene bag. This lower moisture content resulted in lower respiration rate, lower metabolic activity and maintenance of high seed vigour during storage. This results are in similarity with Chattha *et al.*<sup>7</sup> Seeds preserved in ambient

storage in P1 aluminium foil pouch recorded higher germination, seedling length, seedling dry weight, vigour indices compared to P2 high density polythene bag and lower electrical conductivity and moisture content is the indication of lower metabolic activity and respiration rate and maintenance of vigour during the storage. It was observed that under ambient condition loss off seed quality at slow

rate could be due to reduced metabolic activities and inactivation of enzymes required for longer period. Whereas with course of time decline in viability could be due to depletion of food reserves increase fatty acidity, ultra structural changes reduced activity of enzymes and weakening of membrane integrity. These results are in line with the findings of Banovetz and schiener<sup>6</sup> and Diojide<sup>9</sup>.

**Table 1: Analysis of variance for 3 months of storage in Tomato**

S. No.	Characters	Mean squares				Error (d.f=69)
		Treatments (d.f=23)	Factor A (d.f=1)	Factor B (d.f=11)	Factor A X B (d.f=11)	
1.	Germination (%)	6.07	40.36	8.68**	0.36	0.07
2.	Root length (cm)	1.86	3.65**	3.30**	0.25	0.17
3.	Shoot length (cm)	1.301	0.397**	2.689**	0.001**	0.008
4.	Seedling length (cm)	7.12	8.93**	13.94**	0.14	0.20
5.	Seedling dry weight (g)	19.87	0.77	41.30**	0.18	0.35
6.	Vigour Index I	70884	5766	96407**	51280	40103
7.	Vigour Index II	295445	714150**	351842	200985	194358
8.	Electrical conductivity(dsm <sup>-1</sup> )	0.001	0.003	0.003**	0.003**	0.009
9.	Speed of germination	3.597	24.607**	4.885**	0.387**	0.161
10.	Moisture content (%)	0.064	0.003	0.121**	0.013	0.009

\*, \*\* indicate significant at 5% and 1 % level of significance respectively.

**Table 2: Mean performance for 3 months of storage in tomato**

Treatments	Germination (%)			Shoot length(cm)			Root length(cm)			Seedling length(cm)			Seedling dryweight(g)		
	P1	P2	mean	P1	P2	mean	P1	P2	mean	P1	P2	mean	P1	P2	mean
T0	84.25	83.63	83.63	5.80	5.71	5.75	8.05	8.03	8.04	13.67	13.73	13.70	30.55	30.75	30.65
T1	84.25	83.13	83.13	6.67	6.57	7.01	9.13	8.59	8.86	15.79	15.15	15.47	37.49	37.08	37.28
T2	85.50	83.44	83.44	7.12	6.90	6.44	9.32	9.07	9.19	16.34	15.97	16.15	37.82	37.24	37.53
T3	84.88	83.25	83.25	6.50	6.39	6.27	8.52	8.70	8.61	16.02	15.08	15.55	37.13	36.58	36.85
T4	85.00	83.57	83.57	6.34	6.20	7.09	8.49	8.07	8.28	14.97	14.20	14.58	36.78	36.33	36.55
T5	85.63	83.82	83.82	7.16	7.03	6.20	9.35	8.86	9.10	16.28	15.89	16.08	37.45	37.74	37.59
T6	84.50	83.50	83.50	6.27	6.14	7.55	8.25	8.23	8.24	14.83	14.35	14.59	36.24	36.20	36.22
T7	86.50	85.75	85.75	7.64	7.46	7.20	9.64	9.54	9.59	17.45	16.83	17.14	38.78	38.53	38.65
T8	86.00	85.00	85.00	7.30	7.10	6.76	9.56	9.28	9.42	17.18	16.37	16.78	38.58	38.48	38.53
T9	86.00	84.50	84.50	6.79	6.74	6.90	9.04	8.23	8.63	15.88	14.96	15.42	35.93	35.66	35.79
T10	87.5	83.63	83.38	7.00	6.80	7.77	9.19	8.29	8.74	16.18	14.89	15.53	35.56	35.77	35.66
T11	87.75	86.50	86.50	7.77	7.77	7.01	10.02	10.20	10.11	17.97	17.59	17.78	39.80	39.54	39.67
Grand mean	88.97	85.42	84.12	6.86	6.73	6.82	9.04	8.75	8.90	16.04	15.41	15.73	36.84	36.66	36.74
	T	P	T×P	T	P	T×P	T	P	T×P	T	P	T×P	T	P	T×P

SEm±	0.039	0.097	0.028	0.012	0.03	9.12	0.06	0.14	0.04	0.07	0.17	0.04	0.08	0.21	0.06
CD at 5%	0.155	0.187	0.05	0.470	0.057	17.60	0.23	0.27	4.74	0.27	0.32	0.07	0.31	0.405	0.115
Treatments	Vigour index-I			vigour index-II			Speed of germination			Electrical conductivity			Moisture content		
	P1	P2	mean	P1	P2	mean	P1	P2	mean	P1	P2	mean	P1	P2	mean
T0	1377.00	1385.00	1381.00	3105.25	3212.00	3158.63	35.76	33.52	34.64	0.575	0.537	0.556	8.67	8.69	8.68
T1	1486.75	1504.50	1495.63	3470.00	3445.00	3457.50	36.53	35.50	36.01	0.522	0.525	0.524	8.43	8.34	8.38
T2	1555.75	1526.00	1540.88	3531.50	3476.25	3503.88	37.49	36.49	36.99	0.499	0.512	0.506	8.37	8.41	8.39
T3	1451.50	1496.25	1473.88	3448.50	3301.00	3374.75	36.65	35.61	36.13	0.497	0.477	0.487	8.47	8.42	8.44
T4	1457.75	1478.75	1468.25	3443.25	2542.00	2992.63	36.62	35.61	36.11	0.482	0.505	0.494	8.42	8.39	8.40
T5	1562.75	1512.25	1537.50	3557.75	3549.00	3553.38	37.83	36.91	37.37	0.546	0.495	0.521	8.37	8.45	8.41
T6	1110.00	1437.50	1273.75	3339.00	3230.00	3284.50	36.40	35.40	35.90	0.517	0.483	0.500	8.34	8.37	8.35
T7	1623.00	1218.00	1420.50	3657.50	2887.00	3272.25	37.28	36.86	37.07	0.447	0.512	0.480	8.23	8.33	8.28
T8	1645.25	1574.25	1609.75	3621.75	3602.50	3612.13	37.05	36.50	36.77	0.460	0.494	0.477	8.26	8.36	8.31
T9	1377.25	1388.75	1383.00	3283.25	3218.25	3250.75	36.58	35.82	36.20	0.488	0.497	0.493	8.42	8.27	8.35
T10	1385.50	1369.50	1377.50	3236.75	3179.00	3207.88	37.05	35.80	36.42	0.505	0.475	0.490	8.33	8.33	8.33
T11	1686.50	1642.25	1664.38	3728.25	3709.25	3718.75	37.94	36.97	37.45	0.394	0.469	0.432	8.11	8.19	8.15
Grand mean	1476.58	1461.08	1468	3451.90	3279.27	3365	36.93	35.91	36.42	0.494	0.498	0.496	8.37	8.38	8.37
	T	P	T×P	T	P	T×P	T	P	T×P	T	P	T×P	T	P	T×P
SEm±	28.90	70.08	20.40	63.63	155.8	44.90	0.05	0.14	0.04	0.013	0.03	9.68	0.01	0.03	9.68
CD at 5%	115.02	135.25	39.37	253.24	300.69	86.65	0.19	0.27	0.07	0.051	0.057	18.68	0.039	0.057	18.68

### CONCLUSION

It is concluded that from the present investigation that tomato seeds pelletized and treated with the combined treatment of polymer and fungicide T<sub>11</sub>-Vitavax+polymer @6ml/kg seed followed by T<sub>7</sub>.Imidachloprid +polymer@ 4ml/kg recorded significantly higher seed quality parameters. These two seed treatments were found effective in improving the shelf life of seed and it is more beneficial to the farmers. Tomato seeds packed in aluminium foil pouch found more effective for extending seed longevity and maintaining the storability by safe guarding seed from deteriorating.

### Acknowledgement

Author is thankful to Advisor Dr. Abhinav Dayal, Co-Advisor Dr. Prasanth kumar Rai and advisory committee for continuous support, guidance, co-operation. Author feels immense pleasure to thank all the advisory committee members for their constructive

criticism and especially Head and Dean of Post Graduate studies Prof (Dr.) P.W.Ramteke Department of Genetics and Plant Breeding, SHUATS, Allahabad, Uttar Pradesh (U.P), India for providing necessary facilities

### REFERENCES

1. Afzal, R., Mughal, S. M., Munir, M., Sultana, K., Qureshi, R., Arshad, M. and Laghari, M. K., Mycoflora associated with seeds of different sunflower cultivars and its management. *Pakistan Journal of Botany*, **42(1)**: 435-445 (2010).
2. Akter, N., Haque, M. M., Islam, M. R. and Alam, K. M., Seed quality of stored soybean (*Glycine max* L.) as influences by storage containers and storage periods. *The Agriculturists*, **12(1)**: 85-95 (2014).
3. Alam, M. Z., Hamim, I., Ali, M. A. and Ashrafuzzaman, M., Effect of seed treatment on seedling health of chilli. *Journal of Environmental Science and Natural Resources*, **7(1)**: 177-181 (2014).

4. Almeida, A. D. S., Deuner, C., Borges, C. T., Jauer, A., Meneghello, G. E., Tunes, L. M., Villela, F. A. and Zimmer, P. D., Physiological performance of rice seeds treated to thiamethoxam and placed under storage. *American Journal of Plant Sciences*, **5**: 3788-3795. and Biocontrol Agent on Storability of Black Gram (*Vigna mungo L.*). International (2014).
5. Association of Official Seed Analysis-Aosa Seed vigour testing handbook, east lasing AOSA 93p (contribution 32) (1983).
6. Avelar, S. A. G., Souse, F. V. D., Fiss, G., Baudet, L. And Peske, S. T., The use of B. S., Development of polymeric seed coats for Banovetz, S. J. and Schiener, S. M., The effects of seed mass on the seed ecology of *Coreopsis lanceolata*. *Am. Midl.Nat.*, **131(1)**: 65-74 (1994).
7. Chattha, S. H., Ibupoto, K. A., Laghari, M. H., Jamali, L. A. and Baraich, A. A. K., Effect Of Different Packing Materials And Storage Conditions On The Quality Of Wheat Grain. *Pakistan Journal of Agriculture, Engineering and Veterinary Sciences*, **30(2)**: 195-204 (2014).
8. Doijode, S. D., Seed viability and biochemical changes during storage of winter squash (*Cucurbita maxima Duch.*) seeds. *Vegetable Science*. **27**: 168-71 (2000).
9. Doijode, S. D., Seed conservation for preservation of genetic variability in *Salvia splendens*. *J. Orn.Hort.*, **7(3-4)**: 106-10 film coating on the performance of treated corn seed. *Revista Brasileira De Sementes, Hardening and Pelleting Technologies for Rainfed/Garden Land Ecosystems: May 27 Ind. J. Agric. Sci.*, **81(5)**: 423-428 (2004).
10. Jitendra, K., Nisar, K., Kumar, M. B. A., Walia, S., Shakil, N. A., Prasad, R. and Parmer, B. S., Development of polymeric seed coats for seed quality enhancement of Soybean (*Glycine max*). *Indian Journal of Agriculture Sciences*, **77(11)**: 738-43 (2007).
11. Kamara, E. G., Massaquoi, F. B., James, M. S. and George, A., Effects of packing material and treatment on weevil (*Callosobruchus machalatus* (F) Coleoptera: Bruchidae) infestation and quality of cowpea seeds. *African Journal of Agricultural Research*, **9(45)**: 3313-3318 (2014).
12. Krishnasamy, V., Seed pelleting principles and practices. Short Course on Seed Hardening and Pelleting Technologies for Rainfed/Garden Land Ecosystems: May 27 to June 5, *Tamilnadu Agriculture University, Coimbatore, India* (2003).
13. Kunkur, V., Hunje, R., Patil, N. K. B. and Vyakarnhal, B. S., Effect of seed coating with polymer, fungicide and insecticide on seed quality in cotton during storage. *Karnataka Journal of Agriculture Science*, **20(1)**: 137-139 (2007).
14. Kumar, S., Radhamani, J. and Srinivasan, K., Physiological and biochemical changes in seeds of karanj (*Pongamia pinnata*) under different storage conditions. material and treatment on weevil (*Callosobruchus machalatus* (F) Coleoptera: Bruchidae) infestation and quality of cowpea seeds. *African Journal of Agricultural Research*, **9(45)**: 3313-3318 (2011).
15. Monira, U. S., Amin, M. H. A., Aktar, M. M. and Mamun, M. A. A., Effect of packaging materials on seed quality of storage soybean seed. *Bangladesh Research polymer film coat as a micro-container of individual seed facilitates safe storage of N.A.*, Prasad, R. and Parmer, *Journal of Agriculture Sciences*. **77(11)**: 738- 43 *Journal of Plant & Soil Science*, **8(6)**: 1-8 polymeric seed coats for seed quality enhancement of Soybean (*Glycine max*) (2012).
16. Veraja, P. and Rai, P. K., Effect of Polymer Coating, *Chemical Publications Journal*, **7(4)**: 421-427 (2015).
17. Roos, E., Storage behaviour of pelleted, tabeleted taped lettuce seed. *Horticultural science*, **104(2)**: 283-288 (1979).

18. Jacob, S. R., Arun Kumar, M. B., Varghese, E., Sinha S. N., Hydrophilic polymer film coat as a micro-container of individual seed facilitates safe storage of tomato seeds. *Scientia Horticulturae* **204:** 116-122 (2016).
19. Siriwardana, V. M., Abey siriwardena, D. S. D. Z. and Gama-Arachchig, N. S., Effect of seed coatings and storage time on seed quality of rice var.BG 300. Proceedings of the Postgraduate *Institute of Science Research Congress*, Sri Lanka: 9th – 10 th October. 2015; <http://www.researchgate.net/publication/283726407> Soybean (*Glycine max*). *Indian Journal of Agriculture Sciences*, **77(11):** 738-43 (2015).
20. Srinivasan, J., Vijayakumar, A., Srimathi, P., Influence of seed treatment, storage containers and storage periods on storability of the female parent of tomato coth 2. *I.J.S.N.*, **7(3):** 2016: 674-679 (2016).
21. Scott, J. M., Seed coating and treatments and their effects on plant establishment. *Agronomy* **42:** 43-83 (1989).
22. Suma, N. and Srimathi, P., Influence of polymer coating on seed and seedling quality characteristics. *IOSR Journal of Agriculture and Veterinary Science*, **7(5):** 48-50. to June 5, *Tamilnadu Agriculture University*, Coimbatore, India (2014).