

Canopy Temperature Depression and Stay Green: Major Components for Identifying Under Terminal Heat Stress Resistance Genotypes in Wheat

Krishna Kumar Jangid* and Jai Prakash Srivastava

Department of Plant Physiology, Banaras Hindu University, Varanasi, India 221005

*Corresponding Author E-mail: krishnajangid311@gmail.com

Received: 27.05.2018 | Revised: 24.06.2018 | Accepted: 29.06.2018

ABSTRACT

*Analysis of yielding ability in 20 genotypes of spring wheat (*Triticum aestivum*) was done on the basis of morpho-physiological parameter including canopy temperature and stay green trait under terminal heat stress. Effect of terminal heat stress was studied by delayed sowing at three different dates viz., normal (S_1 ; November 26, 2011), late (S_2 ; December 25, 2011) and very late (S_3 ; January 10, 2012) taking 20 genotypes of wheat in randomized block design with three replications. Genotypes showed significant differences in their morpho-physiological traits under different environments. At anthesis stage, chlorophyll content in flag leaf decreased in plant of S_2 and S_3 as compare to S_1 . Canopy temperature increased with advancement in plant growth as well as with delay in sowing. There was significant reduction in total dry matter production, spike number per unit area, spike lets per spike, grains per spike and 1000 grain weight (test weight) with delay in sowing. On the basis of susceptibility index and relative yield loss, genotypic NW-1014 was found to be the most resistant and genotype K-911 the most susceptible to terminal heat stress. NW-1014 was a poor yielder under normal sown condition (S_1) but yielded highest under S_3 , while K-911 yielded highest under S_1 but was poor yielder when sowing was delayed (S_3) among the evaluated genotypes.*

Key words: Chlorophyll content (CC), Canopy temperature (CT), Canopy temperature depression (CTD), Terminal heat stress.

INTRODUCTION

In India, rice-wheat cropping system is prevalent in the northern part. In this region sowing of wheat is delayed due to late harvesting of rice. In such condition wheat crop faces the problem of terminal heat stress and crop yield is limited due to derangement in physiological processes of plant. It is reported

that an increase of 1°C of mean daily temperature during grain filling stage results in reductions in yield by 570 and 620 kg ha⁻¹ in spring and winter wheat cultivars, respectively²³. Adverse effect of increased air temperature are reported on morphological¹³ and physiological²⁴ traits of plants.

Cite this article: Jangid, K.K. and Srivastava, J.P., Canopy Temperature Depression and Stay Green: Major Components for Identifying under Terminal Heat Stress Resistance Genotypes in Wheat, *Int. J. Pure App. Biosci.* 6(3): 374-381 (2018). doi: <http://dx.doi.org/10.18782/2320-7051.6691>

Physiological traits including e.g., maintenance of leaf chlorophyll content for longer duration (stay green characters), lesser reduction in photosynthetic rate, lower canopy temperature, higher leaf conductance are associated with higher terminal heat tolerance in wheat⁶. In wheat genotypes canopy temperature (CT) is a significant parameter to indicate relative tolerance to terminal heat stress and canopy temperature depression (CTD) is observed to be strongly correlated with yield and yield attributes under such stress^{16,4}. Heat stress after anthesis reduces grain weight, quality and total yield by affecting source sink relationship, reducing starch synthesis and limiting movement of photoassimilates to sink or grain⁵. Stay green trait is the important physiological characteristic in which plants show the ability to maintain more photosynthetically active pigments, delayed senescence and thus, facilitate for better growth and yield in plants under drought and heat stress^{1,19}. Determination of canopy temperature depression and stay green traits can increase the efficiency of selecting wheat lines for breeding programme and genetic improvement of crop for terminal heat stress prone regions^{4,14,15}. Stay green trait is associated with higher yield and terminal heat stress resistance in wheat⁹. Selection on the basis of stay green trait is significant in breeding for terminal heat resistance in wheat¹⁰. Higher chlorophyll content is directly correlated to stay green trait and it is determined by genetic information of plant that shows high genetic variability²². Active photosynthetic process governed by green leaves at the grain filling duration is essential for higher grain yield². Decrease in photosynthetic rate under high temperature stress is correlated with chlorophyll loss or early senescence³. It is also proposed that

terminal heat tolerant and susceptible genotypes can be identified on the basis of their susceptibility index⁷.

MATERIAL AND METHODS

Twenty genotypes of wheat viz. K910-4, K612, K910-30, K307, SVPW-1, K512, K607, K911, HUW648, NW-1014, AAI-11, AAI-12, HUW-658, HD-2733, AAI-13, AAI-16, NW-1035, NW-4081, NW-6007 and K-9162 were selected from the “Heat Nursery” of the Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi and sown in randomized block design (2 m × 1.5 m) with three replications. Sowing was done at three different dates for induction of heat stress i.e., November 25, 2011 (normal sowing), December 26, 2011 (late sowing) and January 10, 2012 (very late sowing); henceforth termed as S₁, S₂ and S₃, respectively. Various morpho-physiological characters such as leaf number, days to physiological maturity, chlorophyll content, canopy temperature, canopy temperature depression, grain weight per 30 cm row length (g) were recorded by using random sampling technique. Chlorophyll content was measured by hand held chlorophyll meter “the Minolta SPAD-502” in plant leaves at anthesis and 15 days after anthesis. Measurement of canopy temperature was done by a hand held infrared thermometer-based equipment (Sixth Sense LT-300) by targeting the canopy tissues at an angle of 45° and air temperature was taken by targeting air above 0.5 meter parallel of the particular genotype on in bright sunny days between 1200 and 1400 hrs at anthesis and 15 days after anthesis. Canopy temperature depression was measured by using following formula:

$$\text{CTD} = \text{Air temperature (Ta)} - \text{Canopy temperature (Tc)}$$

Relative yield loss (RYL) was calculated by following formula as given by Fisher and Maurer⁷:

$$\text{RYL} = 1 - [\text{Yd}/\text{Yi}]$$

Where,

YL = Relative yield loss

Yd = Yield under stress condition (S₃)

Yi = Yield under normal condition (S₁)

RESULTS AND DISCUSSION

On perusal of meteorological data, it was observed that reproductive phase of wheat crop experimented gradual increase in ambient temperature as the sowing date was delayed (Table 1). Leaf number per plant was recorded in all the 20 genotype (Fig. 1) at an interval of 7 days starting from 64 days after sowing (DAS) in S_1 , 36 DAS in S_2 and 19 DAS in S_3 . Data indicated that under normal sown condition (S_1) the maximum leaf number in all the genotype was attained at 64 DAS, and then it gradually declined. In all genotypes under S_1 , except in K 910-4, SVPW-1, K-607, AAI-12, HUW-648, and NW-1035, leaves remained green up to 120 DAS. When sowing was done on December 26, 2011, (S_2), the leaf number per plant attained the maximum at about 57 DAS, but leaf senescence occurred at about 105 DAS. When sowing was done on January 10, 2012 (S_3), the maximum leaf number per plant was recorded at about 47 DAS, which gradually decreased and the minimum leaf number was observed at about 82 DAS. Complete senescence of leaves in crops of S_3 sowing was recorded at 82 DAS. Delay in sowing caused reduction in total leaf number per plant and the reduction was severe particularly when sowing was done very late. As sowing was delayed, the duration between attaining maximum leaf number per plant and senescence of all leaves (maturity) reduced significantly. It was observed that when sowing was done at normal date (S_1) crop matured in about 118 days but when it was sown at December 25, and January 10, 2012 crop matured in about 106 and 93 days, respectively. Average grain filling durations under different S_1 , S_2 and S_3 sowing dates were 38, 29 and 24 days, respectively.

Chlorophyll content (SPAD unit) in flag leaves of different genotypes was recorded at anthesis and 15 days after anthesis (Table 2). At anthesis on an average, as compare to S_1 , chlorophyll content in flag leaf of S_2 and S_3 plants decreases marginally. Generally plants of different genotypes under S_2 and S_3 contents relatively lesser amounts of chlorophyll than plants of respective genotype

under S_1 . Chlorophyll content in flag leaf of same genotype declined after 15 days of anthesis and the magnitude of reduction increased progressively with further delay in sowing. Mean chlorophyll content after 15 days after anthesis was the minimum (27.60 SPAD units) in HUW-658 and the maximum (48.80 SPAD units) in K 910-30.

Canopy temperature (CT) ($^{\circ}\text{C}$) was recorded at anthesis and 15 days after anthesis (Table 3). On an average it increased with delay in sowing. While at the anthesis at 15 days after anthesis mean canopy temperatures under S_1 , S_2 and S_3 were 24.03, 25.03 and 29.70 $^{\circ}\text{C}$, respectively, while at 15 days after anthesis CT at these stages increased to 26.56, 28.26 and 38.43 $^{\circ}\text{C}$, respectively. Genotypic differences were evident, as at anthesis stage on an average canopy temperature in different genotype ranged between 27.00 to 32.52 $^{\circ}\text{C}$, while at 15 days after anthesis canopy temperature under S_3 in ranged between 35.40 to 42.20 $^{\circ}\text{C}$.

Canopy temperature depression (CTD) ($^{\circ}\text{C}$) was determined and presented in Table 3. Data were recorded at anthesis as well as at 15 days after anthesis stages. On an average CTD narrowed down with delay in sowing of crop as well as with advancement in plant growth stages. On an average the canopy temperature depression at anthesis stage was the maximum (5.86 $^{\circ}\text{C}$) in genotype AAI-11 and the minimum (3.04 $^{\circ}\text{C}$) in K 910-4, while at 15 days after anthesis CTD was the maximum (3.67 $^{\circ}\text{C}$) in HD 2733 and the minimum (1.76 $^{\circ}\text{C}$) in AAI-13.

As sowing delayed, grain yield 30 cm^{-1} row length decreased accordingly (Table 2). Genotypic differences were significant for S_3 sowing. Under very late sowing condition (S_3) the maximum grain yield (35.22 g 30 cm^{-1} row length) was recorded in NW-1035 and the minimum (16.55 g) in K512. When per cent reduction in grain yield under S_3 as compare to S_1 was calculated, it was observed that the values ranged between 35 to 68%, with the minimum (35.79%) in NW-1035 and the maximum (68%) in K-911 and AAI-11.

On the basis of yield under S_1 the relative yield loss (RLY) in different genotypes under S_3 was calculated (Table 2). The value ranged between 0.243 (NW-1014) to 0.679 (K-911). On the basis of RYL among the studied genotypes, NW-1014 was most resistant to terminal heat stress (minimum RYL), while K-911, the most susceptible (maximum RYL). NW-1014 was a poor yielder under normal sown condition (S_1) but yielded highest under S_3 , while K-911 yielded highest under S_1 but was poor yielder at S_3 among the evaluated genotypes.

It was observed that when sowing was done at three different dates (S_1 , S_2 and S_3) crop matured in about 118, 106 and 93 days respectively. Delay in sowing hastened the senescence of leaf and phases of plant development were completed in shorter durations. It is reported that high temperature hastens plant developmental process (Singh *et al.* 2007). When chlorophyll content in flag leaf of mother shoot was estimated, it was observed that at anthesis under S_1 , S_2 and S_3 differences were not large, but after 15 days of anthesis values were significantly lower in S_3 as compare to S_2 , and in S_2 as compare to S_1 . RYL of different genotypes indicated that genotypes NW-1014 was relatively resistant to terminal heat stress as RYL was of lesser magnitude in this genotype (0.137), while genotype K-911, the most susceptible as RYL was the maximum for this genotype among the studied genotypes. Attempt has been made to identify terminal heat stress resistance and susceptible genotype on the basis of susceptibility index¹². On the basis of relative yield loss, calculated under similar conditions for these genotypes, K911 was found to be most susceptible and NW1014 most stable to terminal heat stress (Table 2). It was evident that genotypic differences were large, and such variations have already been reported by Reynold *et al.*^{14,15}; Joshi *et al.*⁸ and Pandey and Srivastava¹³. It is documented that increased atmospheric drought reduces stomata opening resulting in reduction in transpiration rate and causes warming of leaves as a consequence canopy temperature (CT) increased and

canopy temperature depression (CTD) (air temperature-canopy temperature) narrowed down²¹. In the present study also as the sowing was delayed canopy temperature gradually increased and air temperature to canopy temperature dispersion narrowed down.

When canopy temperature (CT) at anthesis and 15 days after anthesis were correlated (Table 4) with various yield and yield attributes, it was observed that CT at 15 days after anthesis had negative correlation with dry matter accumulation and grain yield 30 cm^{-1} row length. Correlation between 1000 grain weight and canopy temperature at 15 days after anthesis was also negative ($r = -0.364$) (Table 4). These observations indicated that increase in canopy temperature at post anthesis stage decreases dry matter accumulation and seed size in wheat. Canopy temperature at post anthesis stage (15 days after anthesis) may be taken as criteria for selecting terminal heat resistant wheat genotypes. Genotypes with relatively lesser CT at 15 days after anthesis may perform better under late sown condition. Munjal and Rena¹¹ have reported that cool canopy during grain filling period in wheat is an important physiological principle for high temperature stress tolerance. Canopy temperature depression (CTD) has been used as efficient selection criteria for screening wheat genotypes for terminal heat stress resistance¹⁷. Reynolds *et al.*¹⁸ reported that CTD is the most potential assay for heat tolerance of genotypes. Plant growth duration was reduced by delayed in sowing. Significant reduction was observed in grain filling duration. As sowing delayed, crop experienced elevated air temperature during reproductive growth phase. At 15 days after anthesis significant differences were recorded in flag leaf chlorophyll content of crop sown at S_1 , S_2 and S_3 . The chlorophyll content decreased sharply in S_3 as compared to S_1 and in S_2 . Delay in sowing resulted in increased canopy temperature (CT) at anthesis and at 15 days after anthesis. Significant differences were observed in the CT of crops sown at different dates. As the sowing delayed, CT at 15 days after anthesis

increased. Increased canopy temperature at post anthesis (15 days after anthesis) stage decreases dry matter accumulation and seed size in wheat. Maintenance of post anthesis chlorophyll content was directly related with 1000 grain weight under late sown condition. Delay in sowing resulted in increased canopy temperature (CT) at anthesis

and at 15 days after anthesis. Significant differences were observed in the CT of crops sown at different dates. Genotypes with high chlorophyll content and lower canopy temperature at 15 days after anthesis performed better under late sown condition, therefore,




Fig. 1: (A, B,C and D) Variation in leaf number per plant in 20 wheat genotype at an interval of 7 days starting from 64, 71, 78 and 85 days after sowing (DAS) respectively at different dates of sowing (S1 = sowing date November 26, 2011, S2 = December 25, 2011, and S3 = January 10, 2012)

Where (1=K910-4, 2=K612, 3=K910-30, 4=K307, 5=SVPW-1, 6=K512, 7=K607, 8=K911, 9=HUW648, 10=NW-1014, 11=AAI-11, 12=AAI-12, 13=HUW-658, 14=HD-2733, 15=AAI-13, 16=AAI-16, 17= NW-1035, 18=NW-4081, 19=NW-6007 and 20=K-9162aretwenty genotypes of wheat).

Table 1: Weekly meteorological data: Varanasi from November 2011-April 2012

Week no.	Month and date	Rainfall (mm)	Temperature °C			Relative humidity			Sunshine hrs	Evaporation mm
			Max	Min	Difference	Max	Min	Difference		
45	Nov 05-11	0.0	30.3	16.0	14.3	90	34	56	6.9	2.7
46	12-18	0.0	29.6	16.1	13.5	90	41	49	7.4	1.9
47	19-25	0.0	29.9	15.4	14.5	93	43	50	5.9	3.6
48	26-02	0.0	26.1	12.9	13.2	90	41	49	5.9	1.4
49	Dec 03-09	0.0	28.8	14.0	14.8	94	49	45	4.6	1.5
50	10-16	0.0	21.5	10.3	11.2	95	64	31	2.3	1.2
51	17-23	0.0	16.5	7.5	9.0	96	65	31	6.6	1.6
52	24-31	0.0	22.3	7.2	15.1	91	41	50	6.9	2.7
1	Jan 1-7	28.2	20.3	13.0	7.3	92	79	13	3.9	1.7
2	8-14	0.0	18.6	10.6	8.0	89	56	33	3.1	1.8
3	15-21	0.0	22.0	10.6	11.4	86	56	30	6.4	2.8
4	22-28	0.0	21.6	8.3	13.3	86	43	43	7.2	2.9
5	29-04	0.0	22.7	8.5	14.2	88	38	50	8.2	2.5
6	Feb 05-11	0.0	22.7	8.5	14.2	79	48	31	8.5	2.2
7	12-18	0.0	24.3	10.2	14.1	86	51	35	7.8	2.2
8	19-25	0.0	30.3	13.6	16.7	83	41	42	9.7	3.2
9	26-04	0.0	27.8	12.2	15.6	72	32	40	9.8	3.5
10	March 05-11	0.0	29.7	14.5	15.2	74	36	38	8.6	4.0
11	12-18	6.4	28.1	14.5	13.6	75	62	13	7.7	3.8
12	19-25	0.0	33.1	16.2	16.9	68	37	31	9.4	4.5
13	26-01	0.0	36.6	18.1	18.5	59	20	39	9.4	5.5
14	April 02-08	1.3	37.9	21.3	16.6	66	24	42	8.8	4.6
15	09-15	10.3	37.0	21.7	15.3	67	27	40	9.3	5.9
16	16-22	0.0	38.1	22.4	15.7	48	24	24	9.6	8.9
17	23-29	0.0	39.8	23.3	16.5	44	17	27	10.5	7.7

Table 2: Chlorophyll content at anthesis (CC I) and chlorophyll content at 15 days after anthesis (CC II), grain yield (g 30 cm⁻¹ row length), susceptibility index (SI) and relative yield loss (RYL) in 20 genotypes of wheat sown at different dates (S₁ = sowing date November 26, 2011, S₂ = December 25, 2011, and S₃ = January 10, 2012)

S. No.	Genotype	CC I				CC II				Grain yield (g 30 cm ⁻¹)				Relative yield loss
		S ₁	S ₂	S ₃	Mean	S ₁	S ₂	S ₃	Mean	S ₁	S ₂	S ₃	Mean	
1	K910-4	46.70	46.70	48.60	47.33	45.53	43.00	44.00	44.17	55.77	34.91	23.40	55.77	0.580
2	K612	46.30	42.00	38.00	42.10	48.00	45.20	46.00	46.40	55.03	27.65	25.36	55.03	0.539
3	K910-30	48.65	46.70	43.95	46.43	46.97	47.70	48.80	47.82	46.50	43.26	25.02	46.50	0.462
4	K307	46.32	42.90	39.00	42.74	46.93	44.00	42.20	44.37	56.92	22.23	21.09	56.92	0.630
5	SVPW1	48.00	46.44	45.00	46.48	47.00	44.00	39.00	43.33	48.42	34.13	22.01	48.42	0.545
6	K512	44.20	44.30	47.00	45.16	49.20	46.93	37.10	44.41	35.96	35.07	16.55	35.96	0.540
7	K607	48.34	48.60	46.80	47.91	45.60	47.00	39.00	43.86	53.25	39.88	19.85	53.25	0.627
8	K911	47.60	40.40	47.00	45.00	42.35	39.87	36.10	39.44	65.52	40.93	21.03	65.52	0.679**
9	HUW648	48.50	47.00	47.50	47.66	46.10	44.60	44.90	45.20	55.72	33.49	18.87	55.72	0.661
10	NW1014	47.40	45.10	49.00	47.16	46.10	44.70	43.00	44.60	36.75	38.20	27.82	36.75	0.243*
11	AAI1	49.50	46.60	39.00	45.03	48.50	49.50	42.83	46.94	58.86	44.54	19.14	58.86	0.675
12	AAI2	47.20	45.90	47.20	46.76	43.70	42.00	37.40	41.03	57.38	32.02	22.10	57.38	0.615
13	HUW658	47.50	49.40	44.26	47.05	42.35	32.00	27.60	33.98	58.38	30.53	26.72	58.38	0.542
14	HD2733	47.40	46.30	48.00	47.23	48.39	43.67	31.40	41.15	54.64	27.52	29.86	54.64	0.454
15	AAI3	48.00	47.00	46.65	47.21	45.53	43.00	39.00	42.51	55.89	31.57	20.58	55.89	0.632
16	AAI6	41.20	41.20	43.40	41.93	43.00	40.57	36.53	40.03	50.64	38.19	24.35	50.64	0.519
17	NW1035	47.10	46.60	37.00	43.56	40.10	47.00	41.40	42.83	54.85	35.38	35.22	54.85	0.358
18	NW4081	47.80	46.50	48.00	47.43	43.50	46.00	41.00	43.50	54.38	27.97	19.08	54.38	0.649
19	NW6007	45.30	44.40	47.00	45.56	48.00	44.00	42.00	44.66	50.18	32.00	17.42	50.18	0.653
20	K9162	48.00	46.34	44.00	46.11	42.47	44.90	46.00	44.45	43.658	31.23	21.47	43.658	0.508
	MEAN	47.05	45.51	44.81		45.46	43.98	40.26		52.43	34.03	22.84		0.556
ANOVA		SEm±		CD 5%		SEm±		CD 5%		SEm±		CD 5%		
Sowing date (S)		0.14		0.28		0.63		1.25		1.69		3.37		
Genotype (G)		0.37		0.73		1.63		3.25		4.36		8.7		
S x G		0.67		1.34		2.82		5.63		7.56		15.07		

*Most resistant to terminal heatstress, ** Most susceptible to terminal heat stress

Table 3: Canopy temperature at anthesis (CT I) and 15 days after anthesis (CT II), canopy temperature depression at anthesis (CTD I) and 15 days after anthesis (CTD II) in 20 genotypes of wheat sown at different dates (S₁ = sowing date November 26, 2011, S₂ = December 25, 2011, and S₃ = January 10, 2012)

S. No.	Genotype	CT I				CT II				CTD I				CTD II			
		S ₁	S ₂	S ₃	Mean	S ₁	S ₂	S ₃	Mean	S ₁	S ₂	S ₃	Mean	S ₁	S ₂	S ₃	Mean
1	K910-4	24.22	26.22	32.52	27.65	26.65	29.32	35.40	30.45	4.98	3.48	0.68	3.04	2.95	2.28	2.40	2.54
2	K612	23.44	26.22	32.22	27.29	26.22	29.33	37.70	31.08	5.76	3.48	0.98	3.40	3.38	2.27	3.80	3.15
3	K910-30	23.66	25.33	31.22	26.73	28.22	28.22	37.00	31.14	5.54	4.37	1.98	3.96	3.60	3.38	0.80	2.59
4	K307	24.33	25.22	26.22	25.25	27.00	28.65	36.70	30.78	4.87	4.48	6.98	5.44	2.60	2.95	1.10	2.21
5	SVPW1	23.41	25.22	28.43	25.68	26.43	26.43	38.00	30.28	5.79	4.48	4.77	5.01	3.17	5.17	0.60	2.98
6	K512	25.22	26.00	28.00	26.40	26.33	28.00	41.30	31.87	3.98	3.70	5.20	4.29	3.27	1.00	2.80	2.35
7	K607	25.52	25.51	28.66	26.56	26.00	28.22	37.50	30.57	3.68	4.19	4.54	4.13	3.60	3.38	0.30	2.42
8	K911	25.81	26.22	28.41	26.81	29.43	26.34	39.60	31.79	3.39	3.48	4.79	3.88	2.60	5.26	1.80	3.22
9	HUW648	23.91	25.33	31.42	26.88	26.33	28.00	42.20	32.17	5.29	4.37	1.78	3.81	3.27	3.60	1.80	2.89
10	NW1014	24.66	25.33	28.00	25.99	26.22	29.95	36.20	30.79	4.54	4.37	5.20	4.70	3.38	1.65	1.60	2.21
11	AAI1	23.51	24.00	27.00	24.83	26.00	30.00	41.00	32.33	5.69	5.70	6.20	5.86	5.10	0.40	2.80	2.70
12	AAI2	23.41	27.22	31.33	27.32	26.33	28.41	38.30	31.01	5.79	2.48	1.87	3.38	3.27	3.19	0.80	2.42
13	HUW658	23.73	24.00	32.40	26.71	26.00	28.00	40.60	31.53	5.47	5.70	0.80	3.99	3.60	3.60	0.80	2.66
14	HD2733	22.33	24.42	38.00	28.25	25.65	26.33	38.00	29.99	6.87	5.28	1.20	4.45	3.95	5.27	1.80	3.67
15	AAI3	23.66	24.00	27.42	25.02	27.20	30.00	36.50	31.23	5.54	5.70	5.78	5.67	2.40	1.60	1.30	1.76
16	AAI6	23.72	25.00	28.00	25.57	25.00	26.00	39.00	30.00	5.48	4.70	5.20	5.12	4.60	5.60	0.80	3.66
17	NW1035	23.52	25.22	28.50	25.74	26.33	29.66	38.80	31.59	5.68	4.48	4.70	4.95	3.27	1.94	0.40	1.87
18	NW4081	23.52	24.50	28.00	25.34	27.66	28.00	37.00	30.88	5.68	5.20	5.20	5.36	1.94	3.60	0.80	2.11
19	NW6007	24.33	25.33	30.40	26.68	26.22	28.00	38.50	30.90	4.87	4.37	2.80	4.01	3.38	1.80	1.40	2.19
20	K9162	24.81	27.00	28.22	26.67	26.00	29.00	39.40	31.46	4.39	2.70	4.98	4.02	3.60	0.90	2.00	2.16
	MEAN	24.03	25.36	29.70		26.56	28.29	38.43		5.16	4.33	3.78		3.34	2.94	1.49	
ANOVA		SEm±		CD 5%		SEm±		CD 5%		SEm±		CD 5%		SEm±		CD 5%	
Sowing date (S)		1.24		2.48		1.28		2.56		0.06		0.12		0.09		0.19	
Genotype (G)		3.21		6.40		3.32		6.61		0.16		0.32		0.24		0.49	
S x G		5.57		11.09		5.75		11.45		0.28		0.56		0.43		0.86	

Table 4: Correlation coefficients of all the characters under study as obtained under S₃

	Grain yield 30 cm ⁻¹	Harvest index	Spike length	Spikes 30 cm ⁻¹	Grains spike ⁻¹	Grain weight spike ⁻¹	1000 grain weight	Spikelets spike ⁻¹	CC I	CC II	CT I	CT II	CTDI	CTDII
Dry weight 30 cm ⁻¹	0.887	0.237	-0.044	0.681	-0.105	0.019	0.139	0.209	-0.450	-0.002	0.285	-0.376	-0.210	-0.008
Grain yield 30 cm ⁻¹		0.654	-0.141	0.630	-0.182	0.081	0.230	0.009	-0.329	-0.156	0.376	-0.237	-0.269	-0.204
Harvest index			-0.256	0.187	-0.224	0.169	0.322	-0.306	0.012	-0.245	0.324	0.070	-0.231	-0.390
Spike length				0.042	0.635	0.315	-0.147	0.331	-0.479	0.100	-0.276	0.384	0.146	-0.072
Spikes 30 cm ⁻¹					-0.123	-0.342	-0.333	-0.231	-0.368	-0.301	0.362	-0.097	-0.119	0.037
Grains spike ⁻¹						0.401	-0.192	0.192	-0.264	-0.771	-0.255	0.257	0.152	0.054
Grain weight spike ⁻¹							0.772	0.082	-0.085	0.281	-0.140	-0.112	-0.104	-0.023
1000 grain weight								0.011	0.009	0.450	-0.068	-0.364	-0.115	0.001
Spikelets spike ⁻¹									-0.314	0.398	-0.153	-0.167	0.007	-0.061
CC I										-0.238	0.259	-0.137	-0.218	-0.132
CC II											-0.271	-0.245	0.095	0.271
CT I												-0.020	-0.885	0.169
CT II													-0.009	0.207
CTDI														-0.176
CTDII														1.000

REFERENCES

1. Acevedo, E., Nachit, M. and Ortiz Ferrara, G., Effects of heat stress on wheat and possible selection tools for selection in breeding for tolerance, D. A., Saunders, E. d., Wheat for the Nontraditional Warm Areas, the Proceedings of the *International Conference. Mexico, D.F.: CIMMYT: 401-421 (1991).*
2. Al-Khatib, K. and Paulsen, G. M., Photosynthesis and productivity during high-temperature stress of wheat genotypes from major world regions. *Crop Science. 30:* 1127–1132 (1990).
3. Al-khatib, K. and Paulsen, G. M., Mode of high temperature injury to wheat during grain development. *Plant physiology. 61:* 363-368 (1984).
4. Amani I, Fischer, R. A. and Reynolds, M. P., Canopy temperature depression association with yield of irrigated spring wheat cultivars in a hot climate. *J. Agronomy and Crop Science. 176:* 119-129 (1996).
5. Bhullar, S. S. and Jenner, C. F., Differential responses to high temperature of starch and nitrogen accumulation in the grain of four cultivars of wheat. *Australian Journal Plant Physiology 12:* 363–375 (1985).
6. Fischer, R. A., Rees, D., Sayre, K. D., Lu, Z. M., Wheat yield progress is associated with higher stomatal conductance and photosynthesis rate and cooler canopies. *Crop science 38:* 1467-1475 (1998).
7. Fischer, R. A. and Maurer, R., Drought maintenance in spring wheat cultivars, Grain yield response. *Australian Journal Agricultural Research. 29:* 897-912 (1978).
8. Joshi, A. K., Kumari, M., Singh, V. P., Reddy, C. M., Stay green trait: Variation, inheritance and its association with spot blotch resistance in spring wheat (*Triticum aestivum* L.). *Euphytica 153:* 59–71 (2007).
9. Kohli, M. M., Mann, C. and Rajaram, S., Global status and recent progress in breeding wheat for the warmer areas. DA Saunders, ed. Wheat for the nontraditional warmer areas. *Mexico, D.F: CIMMYT pp. 225-241 (1991).*
10. Kumar, U., Joshi, A. K., Kumari, M., Paliwal, R., Identification of QTLs for stay green trait in wheat (*Triticum aestivum* L.) in the ‘Chirya 3’ 3 ‘Sonalika’ population. *Euphytica 174:* 437–445 (2010).
11. Munjal, R., Sharma, V. and Panwar, I. S., Variability parameters, correlation and path coefficients for yield, its components

- and quality traits in bread wheat. *National Journal of Plant Improvement*. **8(2)**: 153-155 (2003).
12. Paliwal, R., Röder, M. S., Kumar, U., Srivastava, J. P. and Joshi, A. K., QTL mapping of terminal heat tolerance in hexaploid wheat (*T. aestivum* L.). *Theoretical and Applied Genetics* **125(3)**: 561-75 (2012).
 13. Pandey, S. K. and Srivastava, J. P., Variability in senescence pattern and membrane stability in wheat genotypes under normal and late sown condition. *Indian Journal of Plant Physiology*. **114**: 169-173 (2009).
 14. Reynolds, M. P., Acevedo, E., Sayre, K. D. and Fisher, R. A., Yield potential in modern wheat varieties : its association with a less competitive ideotype. *Field crops Research* **37**: 149-160 (1994).
 15. Reynolds, M. P., Bolota, M., Delgado, M. I. B., Amani, I., Fischer, R. A., Physiological and morphological traits associated with spring wheat yield under hot, irrigated conditions. *Australian Journal Plant Physiology*. **21**: 717–730 (1994).
 16. Reynolds, M. P., Singh, R. P., Ibrahim, A., Evaluating physiological traits to complement empirical selection for wheat in warm environments. *Euphytica*. **100**: 84-95. (1998).
 17. Reynolds, M. P., Nagarajan, S., Razzaque, M. A. and Ageeb, O. A. A., Using canopy temperature depression to select for yield potential of wheat in heat-stressed environments. *Wheat Special Report No. 42*: Mexico, DF, CIMMYT (1997).
 18. Reynolds, M. P., Nagarajan, S., Razzaque, M. A., Ageeb, O. A. A., Breeding for adaptation to environmental factors: Heat tolerance. In: Reynolds, M. P., Ortiz Monasterio, J. I., McNab, A., (eds) *Application of physiology in wheat breeding* pp. 124–135 (2001).
 19. Rosenow, D. T., Quisenberry, J. E., Wendt, C. W. and Clark, L. E., Drought tolerant sorghum and cotton germplasm. *Agricultural Water Management* **7**: 207–222. (1983).
 20. Singh, R. P., Prasad, P. V. V., Sunita, K., Giri, S. N., Reddy, K. R., Influence of high temperature and breeding for heat tolerance in cotton: a review. *Advances in Agronomy* **93**: 313-385. (2007).
 21. Srivastava, J. P. and Chaturvedi, S. N., Influence of water deficit on transpirational and water relation parameters in wheat. *Annals Arid Zone* **28**: 257-266 (1989).
 22. Thomas, H. and Smart, C. M., Crops that stay green. *Annals of Applied Biology* **123**: 193–201 (1993).
 23. White, J. W. and Reynolds, M. P., A physiological perspective on modelling temperature response in wheat and maize crops. In: White, J. W., (Ed.), *Modelling Temperature Response in Wheat and Maize*, Proceeding of a Workshop. *Cimmyt, El Batan, Mexico*, pp. 8–17 (2001).
 24. Viswanathan, C. and Khanna-Chopra, R., Effect of heat stress on grain growth, starch synthesis and protein synthesis in grains of wheat (*Triticum aestivum* L.) varieties differing in grain weight stability. *Journal of Agronomy and Crop Science*, **186**: 1-7 (2001).