



## Effect of Pre and Post Emergent Herbicides on Growth and Yield Parameter in Transplanted Rice

Bhimashankar<sup>1\*</sup>, Shivaprasad M.<sup>2</sup>, Rajath H. P.<sup>1</sup>, Dinesh Kumar M.<sup>1</sup> and Girijesh G. K.<sup>1</sup>

<sup>1</sup>Department of Agronomy, College of Agriculture, UAHS Shivamogga

<sup>2</sup>Department of Agronomy, ZAHRS, Mudigere, UAHA shivamogga

\*Corresponding Author E-mail: [bhimashankar.rastapur@gmail.com](mailto:bhimashankar.rastapur@gmail.com)

Received: 24.12.2018 | Revised: 30.01.2019 | Accepted: 7.02.2019

### ABSTRACT

A field experiment was conducted during Kharif season 2016 to evaluate the bio-efficacy of new post emergent herbicides along with pre-emergent herbicides in transplanted rice under the hill zone of Karnataka at ZAHRS Mudigere, experiments having ten treatments replicated thrice in RCBD. Among the herbicidal applications, sequential application of Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT followed by (fb) Bispyribac sodium 10 SC @ 20 g a.i ha<sup>-1</sup> at 21 DAT recorded significantly higher growth and yield attributes at harvest viz. plant height (83.10 cm), number of tillers (17.60 hill<sup>-1</sup>), number of green leaves (11.60 hill<sup>-1</sup>), leaf area (169.65 cm<sup>2</sup> hill<sup>-1</sup>), panicle length (18.33 cm), panicle weight (2.89 g panicle<sup>-1</sup>), filled grains (95.60 panicle<sup>-1</sup>), higher grain (5165 kg ha<sup>-1</sup>) and straw yields (6258 kg ha<sup>-1</sup>). Hence, these combinations of herbicides are suitable for the application in transplanted rice. The above treatment was on par with weed free up to 45 DAT (hand weeding) and application of Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb hand weeding at 15 and 30 DAT. Weedy check plots were recorded significantly lower growth and yield attributes which resulted in maximum major nutrients removal by different weeds. However, application of Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Bispyribac sodium 10 SC @ 20 g a.i ha<sup>-1</sup> at 21 DAT was found on par with application of Bensulfuron methyl+Pretilachlor 6.6 GR @ 12.5 kg a.i ha<sup>-1</sup> fb Fenoxyprop p ethyl 9.3 EC @ 1250 ml ha<sup>-1</sup> at 21 DAT and Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Chlorimuron ethyl+Metsulfuron methyl 20 WP @ 8 g a.i ha<sup>-1</sup> at 21 DAT.

**Key words:** Pre and post emergent herbicides, Growth parameter, Yield parameter, Transplanted rice, Hill zone

### INTRODUCTION

Rice is the staple food crop for more than half of the world's population. It is grown in six continents and in more than hundred countries. In Asia, more than two billion people are getting 60-70 per cent of their energy

requirement from rice and its derived products<sup>4</sup>. Human consumption accounts 85 per cent of total production and hence rice deserves a special status among cereals as the world's most important wetland crop.

**Cite this article:** Bhimashankar., Shivaprasad, M., Rajath, H. P., Dinesh Kumar M., and Girijesh, G.K., Effect of Pre and Post Emergent Herbicides on Growth and Yield Parameter in Transplanted Rice, *Int. J. Pure App. Biosci.* 7(1): 435-440 (2019). doi: <http://dx.doi.org/10.18782/2320-7051.7379>

The area under rice in the world is 160.9 m ha with a production of 719.7 mt and average productivity of 4472 kg ha<sup>-1</sup>. In India, rice is grown in an area of 43.97 m ha with an annual production of 106.57 mt and productivity of 2424 kg ha<sup>-1</sup>. In Karnataka, it is grown in 1.33 m ha with a total production of 3.52 mt and the average productivity of 2649 kg ha<sup>-1</sup>.

Rice production is facing various constraints that limit the growth and promoting yield in transplanted rice. Among the various constraints, the weed infestation and shortage of labours are the main problems. The labour requirement during crop establishment is very high at a short span of time, especially during transplanting of seedling and as well as for hand weeding. Hand weeding is a tedious task, time consuming and expensive due to various reasons labour availability is decreasing day by day. Weed infestation is one of the most important factors responsible for yield reduction, which also impairs the quality of the produce due to competition for nutrients, moisture, light and to some extent for space. They harbor many insects, pest and pathogen resulting in poor crop and growth. Losses caused by weeds vary from one location to another, depending on the predominant weed flora and also on the control methods practiced by farmers. In areas where labour is costly and in cases where timely weeding would not be possible due to labour scarcity, herbicide offers an excellent alternative to control weeds in rice fields. In some instances, herbicides offer the most practical, effective and economical means of reducing weed competition and to obtain higher rice yield. Pre-emergent herbicides control the weeds at early stages of crop growth, leaving apart the second flush of the emerged weeds. In order to manage the second flush of weeds, sequential application of pre and post-emergent herbicides may prove promising effective control of weed flora of transplanted rice in absence of manual weed control.

#### MATERIAL AND METHODS

A field experiment was conducted during *Kharif* 2016 at Zonal Agricultural and

Horticultural Research Station, Mudigere. The experiment involves ten treatments includes two pre-emergent herbicides Butachlor 50 EC and Bensulfuron methyl 0.6% G @ 60 g a.i. ha<sup>-1</sup> + Pretilachlor 6% G applied alone as pre-emergent at 3 DAT and six post emergent herbicides 2. 4 D Sodium salt 80, Fenoxypop p ethyl 9.3 EC, Bispyribac sodium 10 SC @, Metsulfuron methyl 20 WP, Chlorimuron ethyl+Metsulfuron methyl 20 WP and Ethoxysulfuron 15 WG. Post emergent herbicide applied as alone and each was preceded by pre emergent herbicides which were compared with weed free up to 45 DAT (hand weeding) and weedy check which was laid out under randomized block design with three replication.

The sowing of cultivar IET-7191 done with a spacing of 20 cm X 10 cm and fertilized with recommended doses of nitrogen, phosphorous and potassium (75:75:90 kg ha<sup>-1</sup>). Five plants were selected randomly in the net plot and labeled for recording the observations in each treatment. Data on growth parameter *viz.*, Plant height, Tillers, No of green leaves, leaf area and total dry matter were recorded at 30, 60, 90 and at harvest. Weed population and weed dry weight are subjected to  $\sqrt{x+0.5}$  transformations by using the Fischer method of ANOVA and the data were analyzed statistically for the test of significance. The level of significance of F test was 5 per cent<sup>5</sup>. The yield attributes and grain yield were recorded at harvesting stage of rice and statistically analyzed.

#### Treatment details:

Details of treatments and the corresponding symbols used during the study are as follows.

T<sub>1</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT followed by (fb) hand weeding at 15 and 30 DAT

T<sub>2</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb 2. 4 D Sodium salt 80 @ 2.5 kg ha<sup>-1</sup> at 21 DAT

T<sub>3</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Fenoxypop p ethyl 9.3 EC 1250 ml ha<sup>-1</sup> at 21 DAT

T<sub>4</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Bispyribac sodium 10 SC @ 20 g a.i ha<sup>-1</sup> at 21 DAT

T<sub>5</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Metsulfuron methyl 20 WP @ 5 g ha<sup>-1</sup> at 21 DAT

T<sub>6</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Ethoxysulfuron 15 WDG @ 15 g ha<sup>-1</sup> at 21 DAT

T<sub>7</sub>: Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Chlorimuron ethyl+Metsulfuron methyl 20 WP @ 8 g ha<sup>-1</sup> at 21 DAT

T<sub>8</sub>: Bensulfuron methyl 0.6 + Pretilachlor 6 GR @ 12.5 kg a.i ha<sup>-1</sup> fb Fenoxypop p ethyl 9.3 EC @ 1250 ml ha<sup>-1</sup> at 21 DAT

T<sub>9</sub>: Weedy check

T<sub>10</sub>: Weed free check up to 45 DAT (hand weeding)

Note: DAT- Days after transplanting, a.i- active ingredient

## RESULTS AND DISCUSSION

**Table 1: Growth parameters of rice as influenced by different weed control treatments in transplanted rice**

Treatment	Plant height (cm)	Number of tillers (hill <sup>-1</sup> )	Number of green leaves (hill <sup>-1</sup> )	Leaf area (cm <sup>2</sup> hill <sup>-1</sup> )	Total dry matter production (g hill <sup>-1</sup> )	Straw yield (kg ha <sup>-1</sup> )
T <sub>1</sub>	83.50	18.20	12.60	186.73	28.87	6,486
T <sub>2</sub>	75.60	14.60	7.80	107.99	23.32	4,995
T <sub>3</sub>	73.30	13.20	6.40	96.09	21.09	4,947
T <sub>4</sub>	83.10	17.60	11.60	169.65	27.81	6,258
T <sub>5</sub>	76.60	15.43	8.23	119.57	23.61	5,314
T <sub>6</sub>	78.00	15.80	9.20	127.37	24.12	5,437
T <sub>7</sub>	79.00	16.20	10.00	140.20	24.86	5,594
T <sub>8</sub>	80.40	16.80	10.80	153.73	25.97	5,769
T <sub>9</sub>	71.45	10.60	4.80	65.50	17.86	3,759
T <sub>10</sub>	85.00	19.00	13.50	205.36	29.61	6,661
S.Em±	1.10	0.60	0.65	13.50	0.83	164.77
C.D. at 5%	3.30	1.80	1.95	40.50	2.49	494.31

**Table 2: Yield parameters of rice as influenced by different weed control treatments in transplanted rice**

Treatment	No. of panicles hill <sup>-1</sup>	Length of Panicle (cm)	Panicle weight (g)	Total no. of grains panicle	Filled grains panicle <sup>-1</sup>	Unfilled grains panicle <sup>-1</sup>	Sterility (%)	Grain yield (kg ha <sup>-1</sup> )
T <sub>1</sub>	17.68	19.31	2.97	107.00	98.80	8.20	7.66	5,441
T <sub>2</sub>	13.66	15.16	2.63	95.03	84.23	10.80	11.34	4,056
T <sub>3</sub>	12.33	13.83	2.49	91.42	79.20	12.22	13.34	3,863
T <sub>4</sub>	17.58	18.33	2.89	104.40	95.60	8.80	8.67	5,165
T <sub>5</sub>	13.80	15.66	2.65	96.66	86.40	10.26	10.77	4,297
T <sub>6</sub>	14.43	17.16	2.67	98.80	88.80	10.00	10.12	4,370
T <sub>7</sub>	15.20	17.32	2.76	100.80	91.20	9.60	9.52	4,511
T <sub>8</sub>	15.55	17.60	2.83	102.60	93.40	9.20	8.95	4,720
T <sub>9</sub>	10.60	10.90	2.18	82.80	68.60	14.20	17.15	2,674
T <sub>10</sub>	19.00	19.66	3.06	110.60	103.00	7.60	6.85	5,647
S.Em±	0.79	0.51	0.07	2.07	2.99	0.41	0.61	172.47
C.D. at 5%	2.36	1.53	0.21	6.21	8.97	1.23	1.86	517.41

The predominant weed flora observed in the experimental field in association with the transplanted rice includes grassy weeds like *Panicum triperon*, *Panicum repens*, *Echinochloa colonum*, sedges like *Cyperus difformis*, *Cyperus procerus*, *Euriocolon sp.* and *Scirpus rayali* and broad leaved weeds like *Monocoria vaginalis*, *Ammania baccifera*, *Rotala verticillaries*, *Eclipta alba*, *Ludwizia parviflora* and *Marselia quadrifolia*. Similar weed species under transplanted rice were also reported by Duary *et al*<sup>3</sup>.

The herbicidal application of Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Bispyribac sodium 10 SC @ 20 g a.i ha<sup>-1</sup> at 21 DAT recorded higher growth and yield parameters like plant height (83.10 cm), number of tillers (17.60 hill<sup>-1</sup>), number of green leaves (11.60 hill<sup>-1</sup>), leaf area (169.65 cm<sup>2</sup> hill<sup>-1</sup>) total dry matter production (27.81 g hill<sup>-1</sup>), number of tillers (17.60 hill<sup>-1</sup>), number of green leaves (11.60 hill<sup>-1</sup>), leaf area (169.65 cm<sup>2</sup> hill<sup>-1</sup>), number of panicle (17.58 hill<sup>-1</sup>), panicle length (18.33 cm), panicle weight (2.89 g panicle<sup>-1</sup>), filled grains (95.60 panicle<sup>-1</sup>), grain yield (5165 kg ha<sup>-1</sup>) and straw yield (6258 kg ha<sup>-1</sup>) at harvest time. This treatment closely followed by application of Butachlor or at 3 DAT fb Chlorimuron ethyl+Metsulfuron methyl 20 WP@ 8 g ha<sup>-1</sup> at 21 DAT and Bensulfuron methyl 0.6 + Pretilachlor 6 GR @ 12.5 kg a.i ha<sup>-1</sup> fb Fenoxypop p ethyl 9.3 EC @ 1250 ml ha<sup>-1</sup> at 21 DAT which are on par with the each other. However, weed free up to 45 DAT recorded highest growth parameters like plant height (85 cm), number of tillers (19.00 hill<sup>-1</sup>), number of green leaves (13.50 hill<sup>-1</sup>), leaf area (205.36 cm<sup>2</sup> hill<sup>-1</sup>) and yield parameters like ), number of panicle (19.00 hill<sup>-1</sup>), panicle length (19.00 cm), panicle weight (3.06 g panicle<sup>-1</sup>), filled grains ( 110.60 panicle<sup>-1</sup>), grain yield (5647 kg ha<sup>-1</sup>) and also straw yield (6661 kg ha<sup>-1</sup>) at harvest time. This was significantly on par with T<sub>4</sub>.

Butachlor found very effective in control weeds at early stages and further different post-emergent weedicides depends on its mode of action and favorable soil condition

effectively controlled the weed flush at later stages. The better weed control and lesser dry weight of the weeds by the sequential application of pre-emergent herbicide (Butachlor) and broad spectrum new post-emergent herbicide such as Bispyribac sodium which controlled total or the most of the weeds very effectively. The rapid absorption of herbicides molecules by the shoots and roots of weeds are responsible for prevention of cell division and inhibition of protein and fat synthesis activity. Similarly the other post-emergent broad spectrum herbicides viz., Ethoxysulfuron, Chlorimuron ethyl+ methyl and Fenoxypop p ethyl controlled the weeds effectively by inhibiting Aceto Lactate Synthase, Aceto Hydroxyl Acid Synthase, acetyl CoA carboxylase enzyme and fatty acid synthesis by activity of both foliage and soil. Sequential application of these herbicides ensured that the reliable control of most of the grasses, sedges and broadleaved weeds throughout the crop growth period as evidenced by lowest weed population and dry weight of the weeds. The results are in conformity with the findings of Parthipan and Ravi<sup>9</sup>, Khaliq *et al.*<sup>6</sup>, and Vaishali Yadava<sup>13</sup>.

During the crop growth period, the number of green leaves and number of tillers per hill increased up to 90 DAT and there after decreases due to senescence. Increase in the growth parameters is mainly attributed to minimum competition of weeds with rice plant for nutrients, moisture and light which showed positive influence on growth components viz., plant height, number of green leaves per hill, number of tillers per hill and finally better dry matter production and its distribution in the crop. Sequential application of pre and above mentioned new post-emergent herbicides resulted in increased dry matter production of plant, which was mainly due to better translocation of photosynthates. This increase in dry matter was a consequence of effective weed control, which reduced the competition of weeds with crop plant and also increased availability of resources. Thus more photosynthates were produced which were transformed into more number of productive tillers per hill, leaf area and leaf area index.

This is due to better weed control efficiency as evidenced by lower weed population and weed dry weight which helped the crop to grow better and absorb more nutrients from the soil. This indicates the replacement of hand weeding by post-emergent application of Bispyribac sodium along with Butachlor pre-emergent application. Further application of Bensulfuron methyl 0.6 +Pretilachlor 6 GR @ 12.5 kg a.i ha<sup>-1</sup> fb Fenoxypop p ethyl 9.3 EC @ 1250 ml ha<sup>-1</sup> at 21 DAT and these two treatment were statistically on par with each other. Similar reasons have been reported by Sreedevi *et al.*<sup>12</sup>, Singh *et al.*<sup>11</sup>, and Vaishali Yadav<sup>13</sup>.

While, statistically least growth parameters *viz.*, plant height (71.44 cm), number of tillers (10.60 hill<sup>-1</sup>), number of green leaves (4.80 hill<sup>-1</sup>), leaf area (65.50 cm<sup>2</sup> hill<sup>-1</sup>) and yield parameter like number of panicle (10.60 hill<sup>-1</sup>), panicle length (10.90 cm), panicle weight (2.18 g panicle<sup>-1</sup>), filled grains (82.80 panicle<sup>-1</sup>), grain yield (2674 kg ha<sup>-1</sup>) and straw yield (3759 kg ha<sup>-1</sup>) was noticed with weedy check. This is due to more competition of the weeds with the crop plants; this is in confirmation with the results of Malaviya *et al.*<sup>8</sup>, and Mallikarjun *et al.*<sup>7</sup>.

### CONCLUSION

The results indicated that among the herbicides tested, sequential application of applied Butachlor 50 EC @ 1.25 kg a.i ha<sup>-1</sup> at 3 DAT fb Bispyribac sodium 10 SC @ 20 g a.i ha<sup>-1</sup> at 21 DAT and Bensulfuron methyl 0.6 + Pretilachlor 6 GR @ 12.5 kg a.i ha<sup>-1</sup> fb Fenoxypop p ethyl 9.3 EC @ 1250 ml ha<sup>-1</sup> at 21 DAT resulted in higher growth and yield parameters like plant height, number of tillers/hill, number of green leaves/hill, leaf area, total dry matter production, number of panicle/hill panicle length, filled grains/panicle, grain and straw yield besides giving broad spectrum of weed control in labour scare area of hill zone of Karnataka.

### REFERENCES

1. Anonymous, Ministry of Agriculture, Govt., of India. <http://www.indiastat.com>. (2015).
2. Anonymous., Agricultural statistics at a glance-2015, Directorate of economics and statistics, Department of Agriculture and Cooperation, New Delhi. (2015b).
3. Duary, B., Mishra, M. M., Dash, R. And Teja, K. C., Weed management in lowland rice. *Indian J. Weed Sci.*, **47(3)**: 224–232 (2015).
4. Geethalakshmi, V., Ramesh, T., Azhagu Palamuthirsolai and Lakshmanan, A., Agronomic evaluation of rice cultivation systems for water and grain productivity. *Archi. Agron. Soil Sci.*, **57(2)**: 159-166 (2011).
5. Gomez, K. A. and Gomez, A. K., *Statistical procedures for Agricultural Research. (2nd Ed.)*. John Wiley and Sons, New York, pp: 105-114 (1984).
6. Khaliq, A., Matloob, A., Mahmood, S., Abbas, R. N. and Khan, M.B., Seeding density and herbicide tank mixtures furnish better weed control and improve growth, yield and quality of direct seeded fine rice. *Int. J. Agric.Bio.*, **12(70)**: 499–508 (2012).
7. Mallikarjun, A. S., Channabasavanna., Sudheendra, S. and Shrinivas. C. S., Effect of herbicides on weed control and yield of wet seeded rice (*Oryza sativa* L.). *The biosci.*, **9(2)**: 581-583 (2014).
8. Malaviya, D. K., Tiwari, R. K., Shukla, V. K., Uma, S. R. and Singh, S. K., *Int. J. Tropical Agric.*, **32(3&4)**: 821-824 (2014).
9. Parthipan, T. AND RAVI, V., Productivity of transplanted rice as influenced by weed control methods. *African J. Agric. Res.*, **11(16)**: 1445-1449 (2016).
10. Singh, M. and Singh, R. P., Influence of crop establishment methods and weed management practices on yield and economics of direct-seeded rice (*Oryza sativa*). *Indian J. Agron.*, **55(3)**: 224-229 (2010).
11. Singh, V., Mangi, L. J., Zahoor A. G., Bhagirath, S. C. and Gupta, R.K., Herbicide options for effective weed management in dry direct seeded rice under scented rice-wheat rotation of

- western Indo-Gangetic Plains. *Crop Protection.*, **81**: 168-176 (2016).
12. Sreedevi, B., Sandhyarani, A., Srinivas, D., Venkatanna, B., Vinaykartheek, A. and Mahender, K. R., Chemical weed control in aerobic rice. *J. Rice Res.*, **9(2)**: 28-31 (2016).
13. Yadav, V., Integrated Weed Management in Aerobic Rice (*Oryza sativa L.*). *M.Sc.(Agri.) Thesis*, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur. (2015).